

Lead Author

Neal Fann U.S. Environmental Protection Agency

Contributing Authors

Terry Brennan Camroden Associates, Inc.

Patrick Dolwick U.S. Environmental Protection Agency Janet L. Gamble U.S. Environmental Protection Agency Vito Ilacqua

U.S. Environmental Protection Agency Laura Kolb

U.S. Environmental Protection Agency **Christopher G. Nolte**

U.S. Environmental Protection Agency **Tanya L. Spero** U.S. Environmental Protection Agency **Lewis Ziska**

U.S. Department of Agriculture

Acknowledgements: Susan Anenberg, U.S. Chemical Safety Board; Amanda Curry Brown, U.S. Environmental Protection Agency; William Fisk, Lawrence Berkeley National Laboratory; Patrick Kinney, Columbia University; Daniel Malashock,* U.S. Department of Health and Human Services, Public Health Service; David Mudarri, CADMUS; Sharon Phillips, U.S. Environmental Protection Agency; Marcus C. Sarofim,* U.S. Environmental Protection Agency;

Recommended Citation: Fann, N., T. Brennan, P. Dolwick, J.L. Gamble, V. Ilacqua, L. Kolb, C.G. Nolte, T.L. Spero, and L. Ziska, 2016: Ch. 3: Air Quality Impacts. *The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment.* U.S. Global Change Research Program, Washington, DC, 69–98. http://dx.doi.org/10.10.7930/J0GQ6VP6

* Chapter Coordinators



Key Findings

Exacerbated Ozone Health Impacts

Key Finding 1: Climate change will make it harder for any given regulatory approach to reduce ground-level ozone pollution in the future as meteorological conditions become increasingly conducive to forming ozone over most of the United States [*Likely, High Confidence*]. Unless offset by additional emissions reductions of ozone precursors, these climate-driven increases in ozone will cause premature deaths, hospital visits, lost school days, and acute respiratory symptoms [*Likely, High Confidence*].

Increased Health Impacts from Wildfires

Key Finding 2: Wildfires emit fine particles and ozone precursors that in turn increase the risk of premature death and adverse chronic and acute cardiovascular and respiratory health outcomes *[Likely, High Confidence]*. Climate change is projected to increase the number and severity of naturally occurring wildfires in parts of the United States, increasing emissions of particulate matter and ozone precursors and resulting in additional adverse health outcomes *[Likely, High Confidence]*.

Worsened Allergy and Asthma Conditions

Key Finding 3: Changes in climate, specifically rising temperatures, altered precipitation patterns, and increasing concentrations of atmospheric carbon dioxide, are expected to contribute to increases in the levels of some airborne allergens and associated increases in asthma episodes and other allergic illnesses *[High Confidence]*.

3.1 Introduction

Changes in the climate affect the air we breathe, both indoors and outdoors. Taken together, changes in the climate affect air quality through three pathways—via outdoor air pollution, aeroallergens, and indoor air pollution. The changing climate has modified weather patterns, which in turn have influenced the levels and location of outdoor air pollutants such as ground-level ozone (O₃) and fine particulate matter.^{1, 2, 3, 4} Increasing carbon dioxide (CO₂) levels also promote the growth of plants that release airborne allergens (aeroallergens). Finally, these changes to outdoor air quality and aeroallergens also affect indoor air quality as both pollutants and aeroallergens infiltrate homes, schools, and other buildings. Climate change influences outdoor air pollutant concentrations in many ways (Figure 1). The climate influences temperatures, cloudiness, humidity, the frequency and intensity of precipitation, and wind patterns,⁵ each of which can influence air quality. At the same time, climate-driven changes in meteorology can also lead to changes in naturally occurring emissions that influence air quality (for example, wildfires, wind-blown dust, and emissions from vegetation). Over longer time scales, human responses to climate change may also affect the amount of energy that humans use, as well as how land is used and where people live. These changes would in turn modify emissions (depending on the fuel source) and thus further influence air quality.^{6, 7} Some air pollutants such as ozone, sulfates, and black carbon also cause changes in

Climate Change and Health—Outdoor Air Quality



Figure 1: This conceptual diagram for an outdoor air quality example illustrates the key pathways by which humans are exposed to health threats from climate drivers, and potential resulting health outcomes (center boxes). These exposure pathways exist within the context of other factors that positively or negatively influence health outcomes (gray side boxes). Key factors that influence vulnerability for individuals are shown in the right box, and include social determinants of health and behavioral choices. Key factors that influence vulnerability at larger scales, such as natural and built environments, governance and management, and institutions, are shown in the left box. All of these influencing factors can affect an individual's or a community's vulnerability through changes in exposure, sensitivity, and adaptive capacity and may also be affected by climate change. See Chapter 1: Introduction for more information.

U.S. Global Change Research Program

climate.⁸ However, this chapter does not consider the climate effects of air pollutants, remaining focused on the health effects resulting from climate-related changes in air pollution exposure.

Poor air quality, whether outdoors or indoors, can negatively affect the human respiratory and cardiovascular systems. Outdoor ground-level ozone and particle pollution can have a range of adverse effects on human health. Current levels of ground-level ozone have been estimated to be responsible for tens of thousands of hospital and emergency room visits, millions of cases of acute respiratory symptoms and school absences, and thousands of premature deaths each year

in the United States.^{9, 10} Fine particle pollution has also been linked to even greater health consequences through harmful cardiovascular and respiratory effects.¹¹

A changing climate can also influence the level of aeroallergens such as pollen, which in turn adversely affect human

health. Rising levels of CO₂ and resulting climate changes alter the production, allergenicity (a measure of how much particular allergens, such as ragweed, affect people), distribution, and seasonal timing of aeroallergens. These changes increase the severity and prevalence of allergic diseases in humans. Higher pollen concentrations and longer pollen seasons can increase allergic sensitization and asthma episodes and thereby limit productivity at work and school.

Finally, climate change may alter the indoor concentrations of pollutants generated outdoors (such as ground-level ozone), particulate matter, and aeroallergens (such as pollen). Changes in the climate may also increase pollutants generated indoors,



Higher pollen concentrations and longer pollen seasons can increase allergic sensitization and asthma episodes.

such as mold and volatile organic compounds. Most of the air people breathe over their lifetimes will be indoors, since people spend the vast majority of their time in indoor environments. Thus, alterations in indoor air pollutant concentrations from climate change have important health implications.

3.2 Climate Impacts on Outdoor Air Pollutants and Health

Changes in the climate affect air pollution levels.^{8, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22} Human-caused climate change has the potential to increase ozone levels,^{1, 4} may have already increased ozone pollution in some regions of the United States,³ and has the potential to affect future concentrations of ozone and fine par-

Human-caused climate change has the potential to increase ozone levels, may have already increased ozone pollution in some regions of the United States, and has the potential to affect future concentrations of ozone and fine particles. ticles (particulate matter smaller than 2.5 microns in diameter, referred to as PM_{2.5}).^{2, 7} Climate change and air quality are both affected by, and influence, several factors; these include the levels and types of pollutants emitted, how land is used, the chemistry governing how these pollutants form in the atmosphere, and weather conditions.

Ground-Level Ozone

Ozone levels and subsequent ozone-related health impacts depend on 1) the amount of pollutants emitted that form ozone, and 2) the meteorological conditions that help determine the amount of ozone produced from those emissions. Both of these factors are expected to change in the future. The emissions of pollutants from anthropogenic (of human origin) sources that form ozone (that is, ozone "precursors") are expected to decrease over the next few decades in the United States.²³ However, irrespective of these changes in emissions, climate change will result in meteorological conditions more favorable to forming ozone. Consequently, attaining national air quality standards for ground-level ozone will also be more difficult, as climate changes offset some of the improvements that would otherwise be expected from emissions reductions. This effect is referred to as the "climate penalty."^{7, 24}

Meteorological conditions influencing ozone levels include air temperatures, humidity, cloud cover, precipitation, wind trajectories, and the amount of vertical mixing in the atmosphere.^{1,} ^{2, 25, 26} Higher temperatures can increase the chemical rates at which ozone is formed and increase ozone precursor emissions from anthropogenic sources and biogenic (vegetative) sources. Lower relative humidity reduces cloud cover and rainfall, promoting the formation of ozone and extending ozone lifetime in the atmosphere. A changing climate will also modify wind patterns across the United States, which will influence local ozone levels. Over much of the country, the worst ozone episodes tend to occur when the local air mass does not change over a period of several days, allowing ozone and ozone precursor emissions

What is Ozone?

Ozone (O₃) is a compound that occurs naturally in Earth's atmosphere but is also formed by human activities. In the stratosphere (10–50 kilometers above the Earth's surface), O₃ prevents harmful solar ultraviolet radiation from reaching the Earth's surface. Near the surface, however, O₃ irritates the respiratory system. Ground-level O₃, a key component of smog, is formed by chemical interactions between sunlight and pollutants including nitrogen oxides (NO_x) and volatile organic compounds (VOCs). The emissions leading to O₃ formation can result from both human sources (for example, motor vehicles and electric power generation) and natural sources (for example, vegetation and wildfires). Occasionally, O₃ that is created naturally in the stratosphere can be mixed downward and contribute to O₃ levels near the surface. Once formed, O₃ can be transported by the wind before eventually being removed from the atmosphere via chemical reactions or by depositing on the surface.

At any given location, O_3 levels are influenced by complex interactions between emissions and meteorological conditions. Generally, higher temperatures, sunnier skies, and lighter winds lead to higher O_3 concentrations by increasing the rate of chemical reactions and by decreasing the extent to which pollutants are mixed with "clean" (less polluted) background air.

For a given level of emissions of O_3 precursors, climate change is generally expected to increase O_3 pollution in the future throughout much of the United States, in part due to higher temperatures and more frequent stagnant air conditions.⁷ Unless offset by additional emissions reductions of ozone precursors, these climate-driven increases in O_3 will cause premature deaths, hospital visits, lost school days, and acute respiratory symptoms.¹⁴

to accumulate over time.^{27, 28} Climate change is already increasing the frequency of these types of stagnation events over parts of the United States,³ and further increases are projected.²⁹ Ozone concentrations near the ground are strongly influenced by upward and downward movement of air ("vertical mixing"). For example, high concentrations of ozone near the ground often occur in urban areas when there is downward movement of air associated with high pressure ("subsidence"), reducing the extent to which locally emitted pollutants are diluted in the atmosphere.³⁰ In addition, high concentrations of ozone can occur in some rural areas resulting from downward transport of ozone from the stratosphere or upper troposphere to the ground.³¹

Aside from the direct meteorological influences, there are also indirect impacts on U.S. ozone levels from other climate-influenced factors. For instance, higher water vapor concentrations due to increased temperatures will increase the natural rate of ozone depletion, particularly in remote areas,³² thus decreasing the baseline level of ozone. Additionally, potential climate-driven increases in nitrogen oxides (NO_x) created by lightning or increased exchange of naturally produced ozone in the stratosphere to the troposphere could also affect ozone in those areas of the country most influenced by background ozone concentrations.³³ Increased occurrences of wildfires due to climate change can also lead to increased ozone concentrations near the ground.³⁴

There is natural year-to-year variability in temperature and other meteorological factors that influence ozone levels.⁷ While global average temperature over 30-year climatic timescales is expected to increase, natural interannual variability will continue to play a significant role in year-to-year changes in temperature.³⁵ Over the next several decades, the influence of climate change on meteorological parameters affecting average levels of ozone is expected to be smaller than the natural interannual variability.³⁶

To address these issues, most assessments of climate impacts on meteorology and associated ozone formation concurrently simulate global and regional chemical transport over multiple years using "coupled" models. This approach can isolate the influence of meteorology in forming ozone from the effect of changes in emissions. The consensus of these model-based assessments is that accelerated rates of photochemical reaction, increased occurrence of stagnation events, and other direct meteorological influences are likely to lead to higher levels of ozone over large portions of the United States.^{8, 14, 16, 17} At the same time, ozone levels in certain regions are projected to decrease as a result of climate change, likely due to localized increases in cloud cover, precipitation, and/or increased dilution resulting from deeper mixed layers. These climate-driven changes in projected ozone vary by season and location, with climate and air quality models showing the most consistency in ozone increases due to climate change in the northeastern United States.8, 37

Generally, ozone levels will likely increase across the United States if ozone precursors are unchanged (see "Research Highlight: Ozone-Related Health Effects" on page 74).^{4, 7, 8} This climate penalty for ozone will offset some of the expected health benefits that would otherwise result from the expected ongoing reductions of ozone precursor emissions, and could prompt the need for adaptive measures (for example, additional ozone precursor emissions reductions) to meet national air quality goals.

Research Highlight: Ozone-Related Health Effects



Los Angeles, California, May 22, 2012. Unless offset by additional emissions reductions of ozone precursors, climate-driven increases in ozone will cause premature deaths, hospital visits, lost school days, and acute respiratory symptoms.

Importance: Ozone is formed in the atmosphere by photochemical reactions of volatile organic compounds (VOCs) and nitrogen oxides (NO_x) in the presence of sunlight. Although U.S. air quality policies are projected to reduce VOC and NO_x emissions,⁵⁶ climate change will increase the frequency of regional weather patterns conducive to increasing ground-level ozone, partially offsetting the expected improvements in air quality.

Objective: Project the number and geographic distribution of additional ozone-related illnesses and premature deaths in the contiguous United States due to climate change between 2000 and 2030 under projected U.S. air quality policies.

Method: Climate scenarios from two global climate models (GCMs) using two different emissions pathways (RCP8.5 and RCP6.0) were dynamically downscaled following Otte et al. (2012)⁵⁷ and used with emissions projections for 2030 and a regional chemical transport model to simulate air quality in the contiguous United States. The resulting changes in ozone in each scenario were then used to compute regional ozone-related health effects attributable to climate change. Ozone-related health impacts were estimated using the environmental Benefits Mapping and Analysis Program–Community Edition (BenMAP–CE). Population exposure was estimated using projected population data from the Integrated Climate and Land Use Scenarios (ICLUS). Further details can be found in Fann et al. (2015).¹⁴

Results: The two downscaled GCM projections result in 1°C to 4°C (1.8°F to 7.2°F) increases in average daily maximum temperatures and 1 to 5 parts per billion increases in daily 8-hour maximum ozone in 2030 throughout the contiguous United States. As seen in previous modeling analyses of climate impacts on ozone, the air quality response to climate change can vary substantially by region and across scenarios.^{22, 58} Unless reductions in ozone precursor emissions offset the influence of climate change, this climate penalty of increased ozone concentrations due to climate change would result in tens to thousands of additional ozone-related illnesses and premature deaths per year.

Research Highlight: Ozone-Related Health Effects, continued

Conclusions: Future climate change will result in higher ozone levels in polluted regions of the contiguous United States. This study isolates the effect of climate change on ozone by using the same emissions of ozone precursors for both 2000-era and 2030-era climate. In addition, this study uses the latest generation of GCM scenarios and represents the most comprehensive analysis of climate-related, ozone-attributable health effects in 2030, and includes not only deaths but also emergency department admissions for asthma, hospital visits for respiratory causes, acute respiratory symptoms, and missed days of school. These results are subject to important uncertainties and limitations. The ozone-climate modeling reflects two scenarios (based on two separate GCMs) considered. Several emissions categories that are important in the formation of ozone and that could be affected by climate, such as motor vehicles, electrical generating units, and wildfires, were left unchanged between the current and future periods. The analysis applied concentration–response relationships from epidemiology studies of historical air pollution episodes; this both implies that the relationship between air pollution and risk will remain constant into the future and that populations will not attempt to reduce their exposure to ozone.

Projected Change in Temperature, Ozone, and Ozone-Related Premature Deaths in 2030



Figure 2. Projected changes in average daily maximum temperature (degrees Fahrenheit), summer average maximum daily 8-hour ozone (parts per billion), and excess ozone-related deaths (incidences per year by county) in the year 2030 relative to the year 2000, following two global climate models and two greenhouse gas concentration pathways, known as Representative Concentration Pathways, or RCPs (see van Vuuren et al. 2011⁴⁹). Each year (2000 and 2030) is represented by 11 years of modeled data for May through September, the traditional ozone season in the United States.

The top panels are based on the National Center for Atmospheric Research/Department of Energy (NCAR/DOE) Community Earth System Model (CESM) following RCP8.5 (a higher greenhouse gas concentration pathway), and the bottom panels are based on the National Aeronautics and Space Administration (NASA) Goddard Institute for Space Studies (GISS) ModelE2-R following RCP6.0 (a moderate greenhouse gas concentration pathway).

The leftmost panels are based on dynamically downscaled regional climate using the NCAR Weather Research and Forecasting (WRF) model, the center panels are based on air quality simulations from the U.S. Environmental Protection Agency (EPA) Community Multiscale Air Quality (CMAQ) model, and the rightmost panels are based on the U.S. EPA Environmental Benefits and Mapping Program (BenMAP).

Fann et al. 2015 reports a range of mortality outcomes based on different methods of computing the mortality effects of ozone changes—the changes in the number of deaths shown in the rightmost panels were computed using the method described in Bell et al. 2004.^{14, 38} (Figure source: adapted from Fann et al. 2015)¹⁴





Projected Change in Ozone-Related Premature Deaths

Figure 3. Projected change in ozone-related premature deaths from 2000 to 2030 by U.S. region and based on CESM/ RCP8.5. Each year (2000 and 2030) is represented by 11 years of modeled data. Ozone-related premature deaths were calculated using the risk coefficient from Bell et al. (2004).³⁸ Boxes indicate 25th, 50th, and 75th percentile change over 11year sample periods, and vertical lines extend to 1.5 times the interquartile range. U.S. regions follow geopolitical boundaries shown in Figure 2. (Figure source: Fann et al. 2015)¹⁴

Air pollution epidemiology studies describe the relationship between a population's historical exposure to air pollutants and the risk of adverse health outcomes. Populations exposed to ozone air pollution are at greater risk of dying prematurely, being admitted to the hospital for respiratory hospital admissions, being admitted to the emergency department, and suffering from aggravated asthma, among other impacts.^{38, 39, 40}

Air pollution health impact assessments combine risk estimates from these epidemiology studies with modeled changes in future or historical air quality changes to estimate the number of air-pollution-related premature deaths and illness.⁴¹ Future ozone-related human health impacts attributable to climate change are projected to lead to hundreds to thousands of premature deaths, hospital admissions, and cases of acute respiratory illnesses per year in the United States in 2030.^{14, 42, 43, 44, 45, 46}

Health outcomes that can be attributed to climate change impacts on air pollution are sensitive to a number of factors noted above—including the climate models used to describe meteorological changes (including precipitation and cloud cover), the models simulating air quality levels (including wildfire incidence), the size and distribution of the population exposed, and the health status of that population (which influences their susceptibility to air pollution; see Ch. 1: Intro-duction).^{42, 47, 48, 49} Moreover, there is emerging evidence that air pollution can interact with climate-related stressors such as

temperature to affect the human physiological response to air pollution.^{39, 42, 50, 51, 52, 53, 54, 55} For example, the risk of dying from exposure to a given level of ozone may increase on warmer days.⁵¹

Particulate Matter

Particulate matter (PM) is a complex mixture of solid- or liquid-phase substances in the atmosphere that arise from both natural and human sources. Principal constituents of PM include sulfate, nitrate, ammonium, organic carbon, elemental carbon, sea salt, and dust. These particles (also known as aerosols) can either be directly emitted or can be formed in the atmosphere from gas-phase precursors. PM smaller than 2.5 microns in diameter (PM_{2.5}) is associated with serious chronic and acute health effects, including lung cancer, chronic obstructive pulmonary disease (COPD), cardiovascular disease, and asthma development and exacerbation.¹¹ The elderly are particularly sensitive to short-term particle exposure, with a higher risk of hospitalization and death.^{59, 60}

As is the case for ozone, atmospheric $PM_{2.5}$ concentrations depend on emissions and on meteorology. Emissions of sulfur dioxide (SO₂), NO_x, and black carbon are projected to decline substantially in the United States over the next few decades due to regulatory controls, ^{56, 61, 62, 63} which will lead to reductions in sulfate and nitrate aerosols.

Climate change is expected to alter several meteorological factors that affect PM_{2.5}, including precipitation patterns and humidity, although there is greater consensus regarding the effects of meteorological changes on ozone than on PM_{2.5}.² Several factors, such as increased humidity, increased stagnation events, and increased biogenic emissions are likely to increase PM_{2.5} levels, while increases in precipitation, enhanced atmospheric mixing, and other factors could decrease PM_{2.5} levels.², 8, 37, 64</sup> Because of the strong influence of changes in precipitation and atmospheric mixing on PM_{2.5} levels, and because there is more variability in projected changes to those variables, there is no consensus yet on whether meteorological changes will lead to a net increase or decrease in PM_{2.5} levels in the United States.², 8, 17, 21, 22, 64, 65

As a result, while it is clear that PM_{2.5} accounts for most of the health burden of outdoor air pollution in the United States,¹⁰ the health effects of climate-induced changes in PM_{2.5} are poorly quantified. Some studies have found that changes in PM_{2.5} will be the dominant driver of air quality-related health effects due to climate change,⁴⁴ while others have suggested a potentially more significant health burden from changes in ozone.⁵⁰

PM resulting from natural sources (such as plants, wildfires, and dust) is sensitive to daily weather patterns, and those fluctuations can affect the intensity of extreme PM episodes (see also Ch. 4: Extreme Events, Section 4.6).⁸ Wildfires are a major source of PM, especially in the western United States during summer.^{66, 67, 68} Because winds carry PM_{2.5} and ozone precursor gases, air pollution from wildfires can affect people even far downwind from the fire location.^{35, 69} PM_{2.5} from wildfires affects human health by increasing the risk of premature death and hospital and emergency department visits.^{70, 71, 72}

Climate change has already led to an increased frequency of large wildfires, as well as longer durations of individual wildfires and longer wildfire seasons in the western United States.⁷³ Future climate change is projected to increase wildfire risks^{74, 75} and associated emissions, with harmful impacts on health.⁷⁶ The area burned by wildfires in North America is expected to increase dramatically over the 21st century due to climate change.^{77, 78} By 2050, changes in wildfires in the western United States are projected to result in 40% increases of organic carbon and 20% increases in elemental carbon aerosol concentrations.⁷⁹ Wildfires may dominate summertime PM_{2.5} concentrations, offsetting even large reductions in anthropogenic PM_{2.5} emissions.²²

Likewise, dust can be an important constituent of PM, especially in the southwest United States. The severity and spatial extent of drought has been projected to increase as a result of climate change,⁸⁰ though the impact of increased aridity on airborne dust PM has not been quantified (see Ch. 4. Extreme Events).²



Nearly 6.8 million children in the United States are affected by asthma, making it a major chronic disease of childhood.

3.3 Climate Impacts on Aeroallergens and Respiratory Diseases

Climate change may alter the production, allergenicity, distribution, and timing of airborne allergens (aeroallergens). These changes contribute to the severity and prevalence of allergic disease in humans. The very young, those with compromised immune systems, and the medically uninsured bear the brunt of asthma and other allergic illnesses. While aeroallergen exposure is not the sole, or even necessarily the most significant factor associated with allergic illnesses, that relationship is part of a complex pathway that links aeroallergen exposure to the prevalence of allergic illnesses, including asthma episodes.^{81,82} On the other hand, climate change may reduce adverse allergic and asthmatic responses in some areas. For example, as some areas become drier, there is the potential for a shortening of the pollen season due to plant stress.

Aeroallergens and Rates of Allergic Diseases in the United States

Aeroallergens are substances present in the air that, once inhaled, stimulate an allergic response in sensitized individuals. Aeroallergens include tree, grass, and weed pollen; indoor and outdoor molds; and other allergenic proteins associated with animal dander, dust mites, and cockroaches.⁸³ Ragweed is the aeroallergen that most commonly affects persons in the United States.⁸⁴

Allergic diseases develop in response to complex and multiple interactions among both genetic and non-genetic factors, including a developing immune system, environmental exposures (such as ambient air pollution or weather conditions), and socioeconomic and demographic factors.^{85, 86, 87} Aeroallergen exposure contributes to the occurrence of asthma episodes, allergic rhinitis or hay fever, sinusitis, conjunctivitis, urticaria (hives), atopic dermatitis or eczema, and anaphylaxis (a severe, whole-body allergic reaction that can be life-threat-

Ragweed Pollen Season Lengthens



Figure 4: Ragweed pollen season length has increased in central North America between 1995 and 2011 by as much as 11 to 27 days in parts of the United States and Canada, in response to rising temperatures. Increases in the length of this allergenic pollen season are correlated with increases in the number of days before the first frost. The largest increases have been observed in northern cities. (Figure source: Melillo et al. 2014. Photo credit: Lewis Ziska, USDA).³⁵

ening).^{84, 88} Allergic illnesses, including hay fever, affect about one-third of the U.S. population, and more than 34 million Americans have been diagnosed with asthma.⁸¹ These diseases have increased in the United States over the past 30 years (see Ch. 1 Introduction). The prevalence of hay fever has increased from 10% of the population in 1970 to 30% in 2000.⁸⁴ Asthma rates have increased from approximately 8 to 55 cases per 1,000 persons to approximately 55 to 90 cases per 1,000 persons over that same time period;⁸⁹ however, there is variation in reports of active cases of asthma as a function of geography and demographics.⁹⁰

Climate Impacts on Aeroallergen Characteristics

Climate change contributes to changes in allergic illnesses as greater concentrations of CO₂, together with higher temperatures and changes in precipitation, extend the start or duration of the growing season, increase the quantity and allergenicity of pollen, and expand the spatial distribution of pollens.^{84, 91, 92, 93, 94}

Historical trends show that climate change has led to changes in the length of the growing season for certain allergenic pollens. For instance, the duration of pollen release for common ragweed (*Ambrosia artemisiifolia*) has been increasing as a function of latitude in recent decades in the midwestern region of North America (see Figure 4). Latitudinal effects on increasing season length were associated primarily with a delay in first frost during the fall season and lengthening of the frost-free period.⁹⁵ Studies in controlled indoor environments find that increases in temperature and CO₂ result in earlier flowering, greater floral numbers, greater pollen production, and increased allergenicity in common ragweed.^{96, 97} In addition, studies using urban areas as proxies for both higher CO₂ and higher temperatures demonstrate earlier flowering of pollen species, which may lead to a longer total pollen season.^{98, 99, 100}

For trees, earlier flowering associated with higher winter and spring temperatures has been observed over a 50-year period

for oak.¹⁰¹ Research on loblolly pine (*Pinus taeda*) also demonstrates that elevated CO_2 could induce earlier and greater seasonal pollen production.¹⁰² Annual birch (*Betula*) pollen production and peak values from 2020 to 2100 are projected to be 1.3 to 2.3 times higher, relative to average values for 2000, with the start and peak dates of pollen release advancing by two to four weeks.¹⁰³

Climate Variability and Effects on Allergic Diseases

Climate change related alterations in local weather patterns, including changes in minimum and maximum temperatures and rainfall, affect the burden of allergic diseases.^{104, 105, 106} The role of weather on the initiation or exacerbation of allergic symptoms in sensitive persons is not well understood.^{86, 107} So-called "thunderstorm asthma" results as allergenic particles are dispersed through osmotic rupture, a phenomenon where cell membranes burst. Pollen grains may, after contact with rain, release part of their cellular contents, including allergen-laced fine particles. Increases in the intensity and frequency of heavy rainfall and storminess over the coming decades is likely to be associated with spikes in aeroallergen concentrations and the potential for related increases in the number and severity of allergic illnesses.^{108, 109}

Potential non-linear interactions between aeroallergens and ambient air pollutants (including ozone, nitrogen dioxide, sulfur dioxide, and fine particulate matter) may increase health risks for people who are simultaneously exposed.^{87, 88, 106, 108, 110, ^{111, 112, 113, 114} In particular, pre-exposure to air pollution (especially ozone or fine particulate matter) may magnify the effects of aeroallergens, as prior damage to airways may increase the permeability of mucous membranes to the penetration of allergens, although existing evidence suggests greater sensitivity but not necessarily a direct link with ozone exposure.¹¹⁵ A recent report noted remaining uncertainties across the epidemiologic, controlled human exposure, and toxicology studies on this emerging topic.³⁹}

3.4 Climate Impacts on Indoor Air Quality and Health: An Emerging Issue

Climate change may worsen existing indoor air problems and create new problems by altering outdoor conditions that affect indoor conditions and by creating more favorable conditions for the growth and spread of pests, infectious agents, and disease vectors that can migrate indoors.¹¹⁶ Climate change can also lead to changes in the mixing of outdoor and indoor air. Reduced mixing of outdoor and indoor air limits penetration of outdoor pollutants into the indoors, but also leads to higher concentrations of pollutants generated indoors since their dilution by outdoor air is decreased.

Indoor air contains a complex mixture of chemical and biological pollutants or contaminants. Contaminants that can be found indoors include carbon monoxide (CO), fine particles (PM_{2.5}), nitrogen dioxide, formaldehyde, radon, mold, and pollen. Indoor air quality varies from building to building and over the course of a day in an individual building.

Public and environmental health professionals have known for decades that poor indoor air quality is associated with adverse respiratory and other health effects.^{116, 117, 118, 119, 120, 121} Since most people spend about 90% of their time indoors,^{122, 123, 124, 125, 126} much of their exposures to airborne pollutants (both those influenced by climate change and those driven by other factors) happen indoors.

Outdoor Air Changes Reflected in Indoor Air

Indoor air pollutants may come from indoor sources or may be transported into the building with outdoor air.^{127, 128} Indoor pollutants of outdoor origin may include ozone, dust, pollen, and fine PM (PM_{2.5}). Even if a building has an outdoor air intake, some air will enter the building through other openings, such as open windows or under doors, or through cracks in the buildings, bypassing any filters and bringing outdoor air pollutants inside.¹²⁹ If there are changes in airborne pollutants of outdoor origin, such as pollen and mold (see Section 3.3) and fine PM from wildfires (see "Particulate Matter" on page 76), there will be changes in indoor exposures to these contaminants. Although indoor fine PM levels from wildfires are typically lower than outdoors (about 50%), because people spend most of their time indoors, most of their exposure to and health effects from wildfire particles (about 80%) will come from particles inhaled



Dampness and mold in U.S. homes are linked to approximately 4.6 million cases of worsened asthma.

Research Highlight: Residential Infiltration and Indoor Air

Importance: Indoor and outdoor air are constantly mixing as air flows through small cracks and openings in buildings (infiltration) in addition to any open doors, windows, and vents. Infiltration or air exchange is driven by differences in barometric pressure, as a result of wind, and of the temperature difference between indoor and outdoor air. The greater this air exchange, the more similar the composition of indoor and outdoor air. Lower air exchange rates accentuate the impact of indoor sources while reducing that of some outdoor pollution. As climate change increases the average temperature of outdoor air, while indoor air continues to be maintained at the same comfortable temperatures, infiltration driven by temperature differences will change as well, modifying exposure to indoor and outdoor air outdoor air pollution sources.

Objective: Project the relative change in infiltration and its effects on exposure to indoor and outdoor air pollution sources for different climates in the United States, between a late-20th century reference and the middle of the current century, in typical detached homes.

Method: The infiltration change projected for 2040–2070 compared to 1970–2000 was modeled for typical single-family residences in urban areas, using temperatures and wind speeds from eight global–regional model combinations for nine U.S. cities (Atlanta, Boston, Chicago, Houston, Los Angeles, Minneapolis, New York, Phoenix, and Seattle). This analysis compares a building to itself, removing the effects of individual building characteristics on infiltration. Indoor temperatures were assumed unchanged between these two periods. Further details can be found in Ilacqua et al. 2015.¹³¹

Results: Because current average yearly temperatures across the contiguous United States are generally below comfortable indoor temperatures, model results indicate that, under future warmer temperatures, infiltration is projected to decrease by about 5%, averaged across cities, seasons, and climate models. Exposure to some pollutants emitted indoors would correspondingly increase, while exposure to some outdoor air pollutants would decrease to some extent. Projections vary, however, among location, seasons, and climate models. In the warmer cities, infiltration during summer months would rise by up to 25% in some models, raising peak exposures to ozone and other related pollutants just when their concentrations are typically highest. Predictions of different models are less consistent for summer months, however, displaying more uncertainty (average modeling relative range of 14%) for summer than for the rest of the year, and in fact not all models predict summer infiltration increases. Modeling uncertainty for the rest of the year is lower than in the summer (relative range less than 6%).

Conclusions: This study shows the potential shifts in residential exposure to indoor and outdoor air pollution sources driven by a changing climate.¹³¹ These conclusions can be applied to small buildings, including single-family homes, row houses, and small offices. Potential adaptations intended to promote energy efficiency by reducing the leakage area of buildings will enhance the effect of decreasing infiltration and increasing exposure to indoor sources. Because of its novelty and lack of additional evidence, the study results should be considered as suggestive of an emerging issue. If replicated by other studies, these findings would add to the evidence on the potential for climate change to alter indoor air quality and further emphasize the impact of indoor air sources on human health. The overall implications of these findings for exposure to ambient and indoor air pollution remain uncertain at present, as they need to be considered along with other determinants of air exchange, such as window-opening behavior, whose relationship with climate change remains poorly characterized.

indoors.¹³⁰ Climate-induced changes in indoor-outdoor temperature differences may somewhat reduce the overall intake of outdoor pollutants into buildings for certain regions and seasons (see "Research Highlight: Residential Infiltration and Indoor Air").¹³¹

Most exposures to high levels of ozone occur outdoors; however, indoor exposures, while lower, occur for much longer time periods. Indoors, ozone concentrations are usually about 10% to 50% of outdoor concentrations; however, since people spend most of their time indoors, most of their exposure to ozone is from indoor air.¹³⁰ Thus, about 45% to 75% of a person's overall exposure to ozone will occur indoors.¹³² About half of the health effects resulting from any outdoor increases in ozone (see Section "Ground-Level Ozone" on page 72) will be due to indoor ozone exposures.¹³⁰ The elderly and children are particularly sensitive to short-term ozone exposure; however, they may spend even more time indoors than the general population and consequently their exposure to ozone is at lower levels for longer periods than the general public.^{133, 134} In addition, ozone entering a building reacts with some organic compounds to produce secondary indoor air pollutants. These reactions lower indoor ozone concentrations but introduce new indoor air contaminants, including other respiratory irritants.¹³⁵

Climate-related increases in droughts and dust storms may result in increases in indoor transmission of dust-borne pathogens, as the dust penetrates the indoor environment. Dust contains particles of biologic origin, including pollen and bacterial and fungal spores. Some of the particles are allergenic.¹³⁶ Pathogenic fungi and bacteria can be found in dust both indoors and outdoors.¹³⁷ For example, in the southwestern United States, spores from the fungi *Coccidiodes*, which can cause valley fever, are found indoors.¹³⁸ The geographic range where *Coccidiodes* is commonly found is increasing. Climate changes, including increases in droughts and temperatures, may be contributing to this spread and to a rise in valley fever (see Ch. 4: Extreme Events).

Legionnaires' disease is primarily contracted from aerosolized water contaminated with Legionella bacteria.139 Legionella bacteria are naturally found outdoors in water and soil; they are also known to contaminate treated water systems in buildings,¹⁴⁰ as well as building cooling systems such as swamp coolers or cooling towers.¹⁴¹ Legionella can also be found indoors inside plumbing fixtures such as showerheads, faucets, and humidifiers.^{142, 143} Legionella can cause outbreaks of a pneumonia known as Legionnaire's disease, which is a potentially fatal infection.¹⁴⁴ Exposure can occur indoors when a spray or mist of contaminated water is inhaled, including mist or spray from showers and swamp coolers.¹⁴⁵ The spread of Legionella bacteria can be affected by regional environmental factors.¹¹⁶ Legionnaires' disease is known to follow a seasonal pattern, with more cases in late summer and autumn, potentially due to warmer and damper conditions.^{146, 147} Cases of Legionnaires' disease are rising in the United States, with an increase of 192% from 2000 to 2009.148, 149 If climate change results in sustained higher temperatures and damper conditions in some areas, there could be increases in the spread and transmission of Legionella.

Contaminants Generated Indoors

Although research directly linking indoor dampness and climate change is not available, information on building science, climate change, and outdoor environmental factors that affect indoor air quality can be used to project how climate change may influence indoor environments.¹³⁰ Climate change could result in increased indoor dampness in at least two ways: 1) if there are more frequent heavy precipitation events and other severe weather events (including high winds, flooding, and winter storms) that result in damage to buildings, allowing water or moisture entry; and 2) if outdoor humidity rises with climate change, indoor humidity and the potential for condensation and dampness will likely rise. Outdoor humidity is usually the largest contributor to indoor dampness on a yearly basis.¹²⁷

Increased indoor dampness and humidity will in turn increase indoor mold, dust mites, bacteria, and other bio-contamination indoors, as well as increase levels of volatile organic compounds (VOCs) and other chemicals resulting from the off-gassing of damp or wet building materials.^{116, 119, 150} Dampness and mold in U.S. homes are linked to approximately 4.6 million cases of worsened asthma and between 8% and 20% of several common respiratory infections, such as acute bronchitis.^{151, 152} If there are climate-induced rises in indoor dampness, there could be increases in adverse health effects related to dampness and mold, such as asthma exacerbation.

Additionally, power outages due to more frequent extreme weather events such as flooding could lead to a number of health effects (see Ch 4: Extreme Events). Heating, ventilation, and air conditioning (HVAC) systems will not function without power; therefore, many buildings could have difficulty maintaining indoor temperatures or humidity. Loss of ventilation, filtration, air circulation, and humidity control can lead to indoor mold growth and increased levels of indoor contaminants,¹⁵³ including VOCs such as formaldehyde.^{119, 154, 155, 156} Power outages are also associated with increases in hospital visits from carbon monoxide (CO) poisoning, primarily due to the incorrect use of backup and portable generators that contaminate indoor air with carbon monoxide.¹³⁵ Following floods, CO poisoning is also associated with the improper indoor use of wood-burning appliances and other combustion appliances designed for use outdoors.¹⁵⁷ There were at least nine deaths from carbon monoxide poisoning related to power outages from 2000 to 2009.¹⁵⁸

Climate factors can influence populations of rodents that produce allergens and can harbor pathogens such as hantaviruses, which can cause Hantavirus Pulmonary Syndrome. Hantaviruses can be spread to people by rodents that infest buildings,¹⁵⁹ and limiting indoor exposure is a key strategy to prevent the spread of hantavirus.¹⁶⁰ Climate change may increase rodent populations in some areas, including indoors, particularly when droughts are followed by periods of heavy rain (see Ch. 4: Extreme Events) and with increases in temperature and rainfall.¹⁶¹ Also, extreme weather events such as heavy rains and flooding may drive some rodents to relocate indoors.¹⁶² Increases in rodent populations may result in increased indoor exposures to rodent allergens and related health effects.^{159, 163, 164} In addition, climate factors may also influence the prevalence of hantaviruses in rodents.^{163, 164} This is a complex dynamic, because climate change may influence rodent populations, ranges, and infection rates.

3.5 Populations of Concern

Certain groups of people may be more susceptible to harm from air pollution due to factors including age, access to healthcare, baseline health status, or other characteristics.⁶⁰ In the contiguous United States, Blacks or African-Americans, women, and the elderly experience the greatest baseline risk from air pollution.¹⁶⁵ The young, older adults, asthmatics, and people whose immune systems are compromised are more vulnerable to indoor air pollutants than the general population.¹⁶⁶ Lower

socioeconomic status and housing disrepair have been associated with higher indoor allergen exposures, though higher-income populations may be more exposed to certain allergens such as dust mites.^{167, 168}

Nearly 6.8 million children in the United States are affected by asthma, making it a major chronic disease of childhood.¹⁶⁹ It is also the main cause of school absenteeism and hospital admissions among children.⁸³ In 2008, 9.3% of American children age 2 to 17 years were reported to have asthma.¹⁶⁹ The onset of asthma in children has been linked to early allergen exposure and viral infections, which act in concert with genetic susceptibility.¹⁷⁰ Children can be particularly susceptible to allergens due to their immature respiratory and immune systems, as well as indoor or outdoor activities that contribute to aeroallergen exposure (see Table 1).^{170, 171, 172, 173}

Minority adults and children also bear a disproportionate burden associated with asthma as measured by emergency department visits, lost work and school days, and overall poorer health status (see Table 1).^{175, 176} Twice as many Black children had asthma-related emergency department visits and hospitalizations compared with White children. Fewer Black and Hispanic children reported using preventative medication like inhaled corticosteroids (ICS) as compared to White children. Black and Hispanic children also had more poorly controlled asthma symptoms, leading to increased emergency department visits and greater use of rescue medications rather than routine daily use of ICS, regardless of symptom control.^{173, 177}

Children living in poverty were 1.75 times more likely to be hospitalized for asthma than their non-poor counterparts. When income is accounted for, no significant difference was observed in the rate of hospital admissions by race or ethnicity. This income effect may be related to access and use of health care and appropriate use of preventive medications such as ICS.¹⁷⁸

Percentage of population with active asthma, by year and selected characteristics: United States, 2001 and 2010.		
Characteristic	Year 2001 %	Year 2010 %
Total	7.3	8.4
Gender		
Male	6.3	7.0
Female	8.3	9.8
Race		
White	7.2	7.8
Black	8.4	11.9
Other	7.2	8.1
Ethnicity		
Hispanic	5.8	7.2
Non-Hispanic	7.6	8.7
Age		
Children (0-17)	8.7	9.3
Adults (18 and older)	6.9	8.2
Age Group		
0-4 years	5.7	6.0
5-14 years	9.9	10.7
15-34 years	8.0	8.6
35-64 years	6.7	8.1
65 years and older	6.0	8.1
Region		
Northeast	8.3	8.8
Midwest	7.5	8.6
South	7.1	8.3
West	6.7	8.3
Federal Poverty Three	shold	
Below 100%	9.9	11.2
100% to < 250%	7.7	8.7
250% to < 450%	6.8	8.2
450% or higher	6.6	7.1
Source: Moorman et al.	2012 ¹⁷⁴	

Table 1: A recent study of children in California found that racial and ethnic minorities are more affected by asthma.¹⁷⁵ Among minority children, the prevalence of asthma varies with the highest rates among Blacks and American Indians/Alaska Natives (17%), followed by non-Hispanic or non-Latino Whites (10%), Hispanics (7%), and Asian Americans (7%).

People with preexisting medical conditions—including hypertension, diabetes, and chronic obstructive pulmonary disorder—are at greater risk for outdoor air pollution-related health effects than the general population.¹⁷⁹ Populations with irregular heartbeats (atrial fibrillation) who were exposed to air pollution and high temperatures experience increased risk.¹⁶⁵ People who live or work in buildings without air conditioning and other ventilation controls or in buildings that are unable to withstand extreme precipitation or flooding events are at greater risk of adverse health effects. Other health risks are related to exposures to poor indoor air quality from mold and other biological contaminants and chemical pollutants emitted from wet building materials. While the presence of air conditioning has been found to greatly reduce the risk of ozone-related deaths, communities with a higher percentage of unemployment and a greater population of Blacks are at greater risk.⁵⁹

3.6 Research Needs

In addition to the emerging issues identified above, the authors highlight the following potential areas for additional scientific and research activity on air quality. Understanding of future air quality and the ability to model future health impacts associated with air quality changes—particularly PM_{2.5} impacts—will be enhanced by improved modeling and projections of climate-dependent variables like wildfires and land-use patterns, as well as improved modeling of ecosystem responses to climate change. Improved collection of data on aeroallergen concentrations in association with other ecosystem variables will facilitate research and modeling of related health impacts.

Future assessments can benefit from research activities that:

- enhance understanding of how interactions among climate-related factors, such as temperature or relative humidity, aeroallergens, and air pollution, affect human health, and how to attribute health impacts to changes in these different risk factors;
- improve the ability to model and project climate change impacts on the formation and fate of air contaminants and quantify the compounded uncertainty in the projections; and
- identify the impacts of changes in indoor dampness, such as mold, other biological contaminants, volatile organic compounds, and indoor air chemistry on indoor air pollutants and health.

Supporting Evidence

PROCESS FOR DEVELOPING CHAPTER

The chapter was developed through technical discussions of relevant evidence and expert deliberation by the report authors at several workshops, teleconferences, and email exchanges. The authors considered inputs and comments submitted by the public, the National Academies of Sciences, and Federal agencies. For additional information on the overall report process, see Appendices 2 and 3.

In addition, the author team held an all-day meeting at the U.S. Environmental Protection Agency National Center for Environmental Assessment in Crystal City, Virginia, on October 15, 2014, to discuss the chapter and develop initial drafts of the Key Findings. A quorum of the authors participated and represented each of the three sections of the chapter—outdoor air quality, aeroallergens, and indoor air quality. These discussions were informed by the results of the literature review as well as the research highlights focused on outdoor air quality and indoor air quality. The team developed Key Finding 2 in response to comments from the National Research Council review panel and the general public.

The Key Findings for outdoor ozone, wildfires, and aeroallergen impacts reflect strong empirical evidence linking changes in climate to these outcomes. When characterizing the human health impacts from outdoor ozone, the team considered the strength of the toxicological, clinical, and epidemiological evidence evaluated in the Ozone Integrated Science Assessment.³⁹ Because there is increasing evidence that climate change will increase the frequency and intensity of wildfire events, this outcome was included as a key finding, despite the inability to quantify this impact with the available tools and data. Because altered patterns of precipitation and increasing levels of CO₂ are anticipated to promote the level of aeroallergens, this outcome is also included as a Key Finding. Finally, because the empirical evidence linking climate change to indoor air quality was more equivocal, we identified this topic as an emerging issue.

KEY FINDING TRACEABLE ACCOUNTS

Exacerbated Ozone Health Impacts

Key Finding 1: Climate change will make it harder for any given regulatory approach to reduce ground-level ozone pollution in the future as meteorological conditions become increasingly conducive to forming ozone over most of the United States [*Likely, High Confidence*]. Unless offset by additional emissions reductions of ozone precursors, these climate-driven increases in ozone will cause premature deaths, hospital visits, lost school days, and acute respiratory symptoms [*Likely, High Confidence*].

Description of evidence base

The Intergovernmental Panel on Climate Change (IPCC) has concluded that warming of the global climate system is unequivocal and that continued increases in greenhouse gas emissions will cause further temperature increases.^{5,} ³⁵ At the same time, there is a well-established relationship between measured temperature and monitored peak ozone levels in the United States.^{1, 25} Numerous climate and air quality modeling studies have also confirmed that increasing temperatures, along with other changes in meteorological variables, are likely to lead to higher peak ozone levels in the United States,^{7, 37} if ozone precursor emissions remain unchanged.

Risk assessments using concentration-response relationships from the epidemiological literature and modeled air quality data have projected substantial health impacts associated with climate-induced changes in air quality. $^{\rm 14,\,42,\,43,\,44,\,46,\,50}$ This literature reports a range of potential changes in ozonerelated, non-accidental mortality due to modeled climate change between the present and 2030 or 2050, depending upon the scenario modeled, the climate and air quality models used, and assumptions about the concentrationresponse function and future populations. Many of the studies suggest that tens to thousands of premature deaths could occur in the future due to climate change impacts on air quality.^{14, 42} At the same time, hundreds of thousands of days of missed school and hundreds of thousands to millions of cases of acute respiratory symptoms also result from the climate-driven ozone increases in the United States.¹⁴

Major uncertainties

Climate projections are driven by greenhouse gas emission scenarios, which vary substantially depending on assumptions for economic growth and climate change mitigation policies. There is significant internal variability in the climate system, which leads to additional uncertainties in climate projections, particularly on a regional basis. Ozone concentrations also depend on emissions that are influenced indirectly by climate change (for example, incidence of wildfires, changes in energy use, energy technology choices), which further compounds the uncertainty. Studies projecting human health impacts apply concentration-response relationships from existing epidemiological studies characterizing historical air quality changes; it is unclear how future changes in the relationship between air quality, population exposure, and baseline health may affect the concentration-response relationship. Finally, these studies do not account for the possibility of a physiological interaction between air pollutants and temperature, which could lead to increases or decreases in air pollution-related deaths and illnesses.

Assessment of confidence and likelihood based on evidence Given the known relationship between temperature and ozone, as well as the numerous air quality modeling studies that suggest climate-driven meteorological changes will yield conditions more favorable for ozone formation in the future, there is **high confidence** that ozone levels will **likely** increase due to climate change, unless offset by reductions in ozone precursor emissions. Based on observed relationships between ozone concentrations and human health responses, there is **high confidence** that any climate-driven increases in ozone will **likely** cause additional cases of premature mortality, as well as increasingly frequent cases of hospital visits and lost school days due to respiratory impacts.

Increased Health Impacts from Wildfires

Key Finding 2: Wildfires emit fine particles and ozone precursors that in turn increase the risk of premature death and adverse chronic and acute cardiovascular and respiratory health outcomes [*Likely, High Confidence*]. Climate change is projected to increase the number and severity of naturally occurring wildfires in parts of the United States, increasing emissions of particulate matter and ozone precursors and resulting in additional adverse health outcomes [*Likely, High Confidence*].

Description of evidence base

The harmful effects of PM concentrations on human health have been well-documented, and there is equally strong evidence linking wildfires to higher PM concentrations regionally. Recent studies have established linkages between wildfire incidence and adverse health outcomes in the nearby population.^{70, 71} Though projections of climate change impacts on precipitation patterns are less certain than those on temperature, there is greater agreement across models that precipitation will decrease in the western United States.⁷⁴ Rising temperatures, decreasing precipitation, and earlier springtime onset of snowmelt are projected to lead to increased frequency and severity of wildfires.^{22, 75, 77}

Major uncertainties

Future climate projections, especially projections of precipitation, are subject to considerable uncertainty. Land management practices, including possible adaptive measures taken to mitigate risk, could alter the frequency and severity of wildfires, the emissions from wildfires, and the associated human exposure to smoke.

Assessment of confidence and likelihood based on evidence

Given the known association between PM and health outcomes and between wildfires and PM concentrations, there is **high confidence** that an increase in wildfire frequency and severity will **likely** lead to an increase in adverse respiratory and cardiac health outcomes. Based on the robustness of the projection by global climate models that precipitation amounts will decrease in parts of the United States, and that summer temperatures will increase, there is **high confidence** that the frequency and severity of wildfire occurrence will **likely** increase, particularly in the western United States.

Worsened Allergy and Asthma Conditions

Key Finding 3: Changes in climate, specifically rising temperatures, altered precipitation patterns, and increasing concentrations of atmospheric carbon dioxide, are expected to contribute to increases in the levels of some airborne allergens and associated increases in asthma episodes and other allergic illnesses [*High Confidence*].

Description of evidence base

There is a large body of evidence supporting the observation that climate change will alter the production, allergenicity, distribution, and timing of aeroallergens. Historical trends show that climate change has led to changes in the length of the growing season for certain allergenic pollens. Climate change also contributes to changes in allergic illnesses as greater concentrations of CO₂, together with higher temperatures and changes in precipitation, extend the start or duration of the growing season, increase the quantity and allergenicity of pollen, and expand the spatial distribution of pollens.^{84, 91, 92, 93, 94} While the role of weather on the initiation or exacerbation of allergic symptoms in sensitive persons is not entirely understood,^{86, 107} increases in intensity and frequency of rainfall and storminess over the coming decades is expected to be associated with spikes in aeroallergen concentrations and the potential for related increases in the number and severity of allergic illnesses.^{108, 109}

These changes in exposure to aeroallergens contribute to the severity and prevalence of allergic disease in humans. Given that aeroallergen exposure is not the sole, or even necessarily the most significant, factor associated with allergic illnesses, that relationship is part of a complex pathway that links exposure to aeroallergens to the prevalence of allergic illnesses.⁸¹ There is consistent and robust evidence that aeroallergen exposure contributes significantly to the occurrence of asthma episodes, hay fever, sinusitis, conjunctivitis, hives, and anaphylaxis.^{84, 88} There is also compelling evidence that allergic diseases develop in response to complex and multiple interactions among both genetic and non-genetic factors, including a developing immune system, environmental exposures (such as ambient air pollution or weather conditions), and socioeconomic and demographic factors.^{85, 86, 87} Finally, there is evidence that potential non-linear interactions between aeroallergens and ambient air pollutants is likely to increase health risks for people who are simultaneously exposed.^{87, 88, 106, 108, 110, 111, 112,} 113, 114

Major uncertainties

The interrelationships between climate variability and change and exposure to aeroallergens are complex. Where

they exist, differences in findings from across the relevant scientific literature may be due to study designs, references to certain species of pollen, geographic characteristics, climate variables, and degree of allergy sensitization.¹⁰⁴ There are also uncertainties with respect to the role of climate change and the extent and nature of its effects as they contribute to aeroallergen-related diseases, especially asthma.⁹¹ Existing uncertainties can be addressed through the development of standardized approaches for measuring exposures and tracking outcomes across a range of allergic illnesses, vulnerable populations, and geographic proximity to exposures.⁸²

Assessment of confidence and likelihood based on evidence

The scientific literature suggests that there is **high confidence** that changes in climate, including rising temperatures and altered precipitation patterns, will affect the concentration, allergenicity, season length, and spatial distribution of a number of aeroallergens, and these changes are expected to impact the prevalence of some allergic diseases, including asthma attacks.

DOCUMENTING UNCERTAINTY

This assessment relies on two metrics to communicate the degree of certainty in Key Findings. See Appendix 4: Documenting Uncertainty for more on assessments of likelihood and confidence.

Confidence Level
Very High
Strong evidence (established theory, multiple sources, consistent results, well documented and accepted methods, etc.), high consensus
High
Moderate evidence (several sourc- es, some consistency, methods vary and/or documentation limited, etc.), medium consensus
Medium
Suggestive evidence (a few sourc- es, limited consistency, models incomplete, methods emerging, etc.), competing schools of thought
Low
Inconclusive evidence (limited sources, extrapolations, inconsis- tent findings, poor documentation and/or methods not tested, etc.)

Likelihood		
Very Likely		
≥ 9 in 10		
Likely		
≥ 2 in 3		
As Likely As Not		
\approx 1 in 2		
Unlikely		
\leq 1 in 3		
Very Unlikely		
≤ 1 in 10		

PHOTO CREDITS

- Pg. 69 –Girl with inhaler : © Stephen Welstead/LWA/Corbis
- Pg. 70–Firefighters walking in smoke: © Ted Soqui/Corbis
- Pg. 72-Ragweed pollen: Courtesy of Roy Morsch/Corbis
- Pg. 74–L.A. smog: © Ringo Chiu/ZUMA Press/Corbis
- Pg. 77 –Girl with inhaler : © Stephen Welstead/LWA/Corbis
- Pg. 79–Moldy archway: Courtesy of Bart Everson/flickr

References

- Bloomer, B.J., J.W. Stehr, C.A. Piety, R.J. Salawitch, and R.R. Dickerson, 2009: Observed relationships of ozone air pollution with temperature and emissions. *Geophysical Research Letters*, 36, L09803. http://dx.doi. org/10.1029/2009gl037308
- Dawson, J.P., B.J. Bloomer, D.A. Winner, and C.P. Weaver, 2014: Understanding the meteorological drivers of U.S. particulate matter concentrations in a changing climate. *Bulletin* of the American Meteorological Society, 95, 521-532. http:// dx.doi.org/10.1175/BAMS-D-12-00181.1
- Leibensperger, E.M., L.J. Mickley, and D.J. Jacob, 2008: Sensitivity of US air quality to mid-latitude cyclone frequency and implications of 1980–2006 climate change. *Atmospheric Chemistry and Physics*, 8, 7075-7086. http://dx.doi. org/10.5194/acp-8-7075-2008
- 4. Weaver, C.P., E. Cooter, R. Gilliam, A. Gilliland, A. Grambsch, D. Grano, B. Hemming, S.W. Hunt, C. Nolte, D.A. Winner, X.-Z. Liang, J. Zhu, M. Caughey, K. Kunkel, J.-T. Lin, Z. Tao, A. Williams, D.J. Wuebbles, P.J. Adams, J.P. Dawson, P. Amar, S. He, J. Avise, J. Chen, R.C. Cohen, A.H. Goldstein, R.A. Harley, A.L. Steiner, S. Tonse, A. Guenther, J.-F. Lamarque, C. Wiedinmyer, W.I. Gustafson, L.R. Leung, C. Hogrefe, H.-C. Huang, D.J. Jacob, L.J. Mickley, S. Wu, P.L. Kinney, B. Lamb, N.K. Larkin, D. McKenzie, K.-J. Liao, K. Manomaiphiboon, A.G. Russell, E. Tagaris, B.H. Lynn, C. Mass, E. Salathé, S.M. O'Neill, S.N. Pandis, P.N. Racherla, C. Rosenzweig, and J.-H. Woo, 2009: A preliminary synthesis of modeled climate change impacts on U.S. regional ozone concentrations. Bulletin of the American Meteorological Society, 90, 1843-1863. http:// dx.doi.org/10.1175/2009BAMS2568.1
- IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley (Eds.), 1535 pp. Cambridge University Press, Cambridge, UK and New York, NY. http://www.climatechange2013.org
- Bernard, S.M., J.M. Samet, A. Grambsch, K.L. Ebi, and I. Romieu, 2001: The potential impacts of climate variability and change on air pollution-related health effects in the United States. *Environmental Health Perspectives*, 109, 199-209. PMC1240667
- Jacob, D.J. and D.A. Winner, 2009: Effect of climate change on air quality. *Atmospheric Environment*, 43, 51-63. http:// dx.doi.org/10.1016/j.atmosenv.2008.09.051

- Fiore, A.M., V. Naik, D.V. Spracklen, A. Steiner, N. Unger, M. Prather, D. Bergmann, P.J. Cameron-Smith, I. Cionni, W.J. Collins, S. Dalsøren, V. Eyring, G.A. Folberth, P. Ginoux, L.W. Horowitz, B. Josse, J.-F. Lamarque, I.A. MacKenzie, T. Nagashima, F.M. O'Connor, M. Righi, S.T. Rumbold, D.T. Shindell, R.B. Skeie, K. Sudo, S. Szopa, T. Takemura, and G. Zeng, 2012: Global air quality and climate. *Chemical Society Reviews*, 41, 6663-6683. http://dx. doi.org/10.1039/C2CS35095E
- Anenberg, S.C., I.J. West, A.M. Fiore, D.A. Jaffe, M.J. Prather, D. Bergmann, K. Cuvelier, F.J. Dentener, B.N. Duncan, M. Gauss, P. Hess, J.E. Jonson, A. Lupu, I.A. Mackenzie, E. Marmer, R.J. Park, M.G. Sanderson, M. Schultz, D.T. Shindell, S. Szopa, M.G. Vivanco, O. Wild, and G. Zeng, 2009: Intercontinental impacts of ozone pollution on human mortality. *Environmental Science & Technology*, 43, 6482-6287. http://dx.doi.org/10.1021/es900518z
- Fann, N., A.D. Lamson, S.C. Anenberg, K. Wesson, D. Risley, and B.J. Hubbell, 2012: Estimating the national public health burden associated with exposure to ambient PM2.5 and ozone. *Risk Analysis*, **32**, 81-95. http://dx.doi. org/10.1111/j.1539-6924.2011.01630.x
- EPA, 2009: Integrated Science Assessment for Particulate Matter. EPA/600/R-08/139F. National Center for Environmental Assessment, Office of Research and Development, U.S. Environmental Protection Agency, Research Triangle Park, NC. http://cfpub.epa.gov/ncea/cfm/recordisplay. cfm?deid=216546
- Chang, H.H., J. Zhou, and M. Fuentes, 2010: Impact of climate change on ambient ozone level and mortality in south-eastern United States. *International Journal of Environmental Research and Public Health*, 7, 2866-2880. http://dx.doi.org/10.3390/ijerph7072866
- Chang, H.H., H. Hao, and S.E. Sarnat, 2014: A statistical modeling framework for projecting future ambient ozone and its health impact due to climate change. *Atmospheric Environment*, **89**, 290-297. http://dx.doi.org/10.1016/j. atmosenv.2014.02.037
- Fann, N., C.G. Nolte, P. Dolwick, T.L. Spero, A. Curry Brown, S. Phillips, and S. Anenberg, 2015: The geographic distribution and economic value of climate change-related ozone health impacts in the United States in 2030. *Journal* of the Air & Waste Management Association, 65, 570-580. http://dx.doi.org/10.1080/10962247.2014.996270

- Jackson, J.E., M.G. Yost, C. Karr, C. Fitzpatrick, B.K. Lamb, S.H. Chung, J. Chen, J. Avise, R.A. Rosenblatt, and R.A. Fenske, 2010: Public health impacts of climate change in Washington State: Projected mortality risks due to heat events and air pollution. *Climatic Change*, **102**, 159-186. http://dx.doi.org/10.1007/s10584-010-9852-3
- Kelly, J., P.A. Makar, and D.A. Plummer, 2012: Projections of mid-century summer air-quality for North America: Effects of changes in climate and precursor emissions. *Atmospheric Chemistry and Physics*, **12**, 5367-5390. http://dx.doi. org/10.5194/acp-12-5367-2012
- Penrod, A., Y. Zhang, K. Wang, S.-Y. Wu, and L.R. Leung, 2014: Impacts of future climate and emission changes on U.S. air quality. *Atmospheric Environment*, **89**, 533-547. http://dx.doi.org/10.1016/j.atmosenv.2014.01.001
- Pfister, G.G., S. Walters, J.F. Lamarque, J. Fast, M.C. Barth, J. Wong, J. Done, G. Holland, and C.L. Bruyère, 2014: Projections of future summertime ozone over the U.S. *Journal of Geophysical Research: Atmospheres*, **119**, 5559-5582. http:// dx.doi.org/10.1002/2013JD020932
- Sheffield, P.E., J.L. Carr, P.L. Kinney, and K. Knowlton, 2011: Modeling of regional climate change effects on ground-level ozone and childhood asthma. *American Journal of Preventive Medicine*, 41, 251-257. http://dx.doi.org/10.1016/j.amepre.2011.04.017
- Sheffield, P.E., K.R. Weinberger, and P.L. Kinney, 2011: Climate change, aeroallergens, and pediatric allergic disease. *Mount Sinai Journal of Medicine*, 78, 78-84. http://dx.doi. org/10.1002/msj.20232
- 21. Tai, A.P.K., L.J. Mickley, and D.J. Jacob, 2012: Impact of 2000–2050 climate change on fine particulate matter (PM2.5) air quality inferred from a multi-model analysis of meteorological modes. *Atmospheric Chemistry and Physics*, **12**, 11329-11337. http://dx.doi.org/10.5194/acp-12-11329-2012
- 22. Val Martin, M., C.L. Heald, J.F. Lamarque, S. Tilmes, L.K. Emmons, and B.A. Schichtel, 2015: How emissions, climate, and land use change will impact mid-century air quality over the United States: A focus on effects at National Parks. *Atmospheric Chemistry and Physics*, **15**, 2805-2823. http://dx.doi.org/10.5194/acp-15-2805-2015

- 23. EPA, 2014: Regulatory Impact Analysis of the Proposed Revisions to the National Ambient Air Quality Standards for Ground-Level Ozone. EPA-452/P-14-006. U.S. Environmental Protection Agency, Office of Air and Radiation, Office of Air Quality Planning and Standards, Research Triangle Park, NC. http://epa.gov/ttn/ecas/regdata/ RIAs/20141125ria.pdf
- Wu, S., L.J. Mickley, E.M. Leibensperger, D.J. Jacob, D. Rind, and D.G. Streets, 2008: Effects of 2000–2050 global change on ozone air quality in the United States. *Journal of Geophysical Research: Atmospheres*, 113, D06302. http://dx. doi.org/10.1029/2007JD008917
- 25. Camalier, L., W. Cox, and P. Dolwick, 2007: The effects of meteorology on ozone in urban areas and their use in assessing ozone trends. *Atmospheric Environment*, **41**, 7127-7137. http://dx.doi.org/10.1016/j.atmosenv.2007.04.061
- Davis, J., W. Cox, A. Reff, and P. Dolwick, 2011: A comparison of CMAQ-based and observation-based statistical models relating ozone to meteorological parameters. *Atmospheric Environment*, 45, 3481-3487. http://dx.doi.org/10.1016/j. atmosenv.2010.12.060
- 27. Leung, L.R. and W.I. Gustafson, Jr., 2005: Potential regional climate change and implications to U.S. air quality. *Geophysical Research Letters*, **32**, L16711. http://dx.doi.org/10.1029/2005GL022911
- Zhu, J. and X.-Z. Liang, 2013: Impacts of the Bermuda high on regional climate and ozone over the United states. *Journal of Climate*, 26, 1018-1032. http://dx.doi.org/10.1175/ JCLI-D-12-00168.1
- Horton, D.E., Harshvardhan, and N.S. Diffenbaugh, 2012: Response of air stagnation frequency to anthropogenically enhanced radiative forcing. *Environmental Research Letters*, 7, 044034. http://dx.doi.org/10.1088/1748-9326/7/4/044034
- Haman, C.L., E. Couzo, J.H. Flynn, W. Vizuete, B. Heffron, and B.L. Lefer, 2014: Relationship between boundary layer heights and growth rates with ground-level ozone in Houston, Texas. *Journal of Geophysical Research: Atmospheres*, 119, 6230-6245. http://dx.doi.org/10.1002/2013JD020473
- Zhang, L., D.J. Jacob, X. Yue, N.V. Downey, D.A. Wood, and D. Blewitt, 2014: Sources contributing to background surface ozone in the US Intermountain West. *Atmospheric Chemistry and Physics*, 14, 5295-5309. http://dx.doi. org/10.5194/acp-14-5295-2014

- Murazaki, K. and P. Hess, 2006: How does climate change contribute to surface ozone change over the United States? *Journal of Geophysical Research - Atmospheres*, 111, D05301. http://dx.doi.org/10.1029/2005JD005873
- 33. Kirtman, B., S.B. Power, J.A. Adedoyin, G.J. Boer, R. Bojariu, I. Camilloni, F.J. Doblas-Reyes, A.M. Fiore, M. Kimoto, G.A. Meehl, M. Prather, A. Sarr, C. Schär, R. Sutton, G.J. van Oldenborgh, G. Vecchi, and H.J. Wang, 2013: Nearterm climate change: Projections and predictability. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley, Eds. Cambridge University Press, Cambridge, UK and New York, NY, USA, 953–1028. http://dx.doi.org/10.1017/CBO9781107415324.023
- Stavros, E.N., D. McKenzie, and N. Larkin, 2014: The climate-wildfire-air quality system: Interactions and feedbacks across spatial and temporal scales. *Wiley Interdisciplinary Reviews: Climate Change*, 5, 719-733. http://dx.doi. org/10.1002/wcc.303
- Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds., 2014: *Climate Change Impacts in the United States: The Third National Climate Assessment*. U.S. Global Change Research Program, Washington, D.C., 842 pp. http://dx.doi. org/10.7930/J0Z31WJ2
- 36. Rieder, H.E., A.M. Fiore, L.W. Horowitz, and V. Naik, 2015: Projecting policy-relevant metrics for high summertime ozone pollution events over the eastern United States due to climate and emission changes during the 21st century. *Journal of Geophysical Research: Atmospheres*, **120**, 784-800. http://dx.doi.org/10.1002/2014JD022303
- Fiore, A.M., V. Naik, and E.M. Leibensperger, 2015: Air quality and climate connections. *Journal of the Air & Waste Management Association*, 65, 645-686. http://dx.doi.org/10 .1080/10962247.2015.1040526
- Bell, M.L., A. McDermott, S.L. Zeger, J.M. Samet, and F. Dominici, 2004: Ozone and short-term mortality in 95 US urban communities, 1987-2000. *JAMA: The Journal of the American Medical Association*, **292**, 2372-2378. http://dx. doi.org/10.1001/jama.292.19.2372
- EPA, 2013: Integrated Science Assessment for Ozone and Related Photochemical Oxidants. EPA 600/R-10/076F, 1251 pp. U.S. Environmental Protection Agency, National Center for Environmental Assessment, Office of Research and Development, Research Triangle Park, NC. http:// cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=247492

- Jerrett, M., R.T. Burnett, C.A. Pope, K. Ito, G. Thurston, D. Krewski, Y. Shi, E. Calle, and M. Thun, 2009: Long-term ozone exposure and mortality. *The New England Journal of Medicine*, 360, 1085-1095. http://dx.doi.org/10.1056/ NEJMoa0803894
- Hubbell, B., N. Fann, and J. Levy, 2009: Methodological considerations in developing local-scale health impact assessments: Balancing national, regional, and local data. *Air Quality, Atmosphere & Health*, 2, 99-110. http://dx.doi. org/10.1007/s11869-009-0037-z
- Post, E.S., A. Grambsch, C. Weaver, P. Morefield, J. Huang, L.-Y. Leung, C.G. Nolte, P. Adams, X.-Z. Liang, J.-H. Zhu, and H. Mahone, 2012: Variation in estimated ozone-related health impacts of climate change due to modeling choices and assumptions. *Environmental Health Perspectives*, **120**, 1559-1564. http://dx.doi.org/10.1289/ehp.1104271
- Selin, N.E., S. Wu, K.M. Nam, J.M. Reilly, S. Paltsev, R.G. Prinn, and M.D. Webster, 2009: Global health and economic impacts of future ozone pollution. *Environmental Research Letters*, 4, 044014. http://dx.doi.org/10.1088/1748-9326/4/4/044014
- Tagaris, E., K.-J. Liao, A.J. DeLucia, L. Deck, P. Amar, and A.G. Russell, 2009: Potential impact of climate change on air pollution-related human health effects. *Environmental Science & Technology*, 43, 4979-4988. http://dx.doi. org/10.1021/es803650w
- Garcia-Menendez, F., R.K. Saari, E. Monier, and N.E. Selin, 2015: U.S. air quality and health benefits from avoided climate change under greenhouse gas mitigation. *Environmental Science & Technology*, 49, 7580-7588. http://dx.doi. org/10.1021/acs.est.5b01324
- Bell, M.L., R. Goldberg, C. Hogrefe, P.L. Kinney, K. Knowlton, B. Lynn, J. Rosenthal, C. Rosenzweig, and J.A. Patz, 2007: Climate change, ambient ozone, and health in 50 US cities. *Climatic Change*, 82, 61-76. http://dx.doi.org/10.1007/s10584-006-9166-7
- Riahi, K., S. Rao, V. Krey, C. Cho, V. Chirkov, G. Fischer, G. Kindermann, N. Nakicenovic, and P. Rafaj, 2011: RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Climatic Change*, **109**, 33-57. http://dx.doi. org/10.1007/s10584-011-0149-y
- Taylor, K.E., R.J. Stouffer, and G.A. Meehl, 2012: An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93, 485-498. http://dx.doi.org/10.1175/BAMS-D-11-00094.1

- van Vuuren, D.P., J. Edmonds, M. Kainuma, K. Riahi, A. Thomson, K. Hibbard, G.C. Hurtt, T. Kram, V. Krey, J.-F. Lamarque, T. Masui, M. Meinshausen, N. Nakicenovic, S.J. Smith, and S.K. Rose, 2011: The representative concentration pathways: An overview. *Climatic Change*, **109**, 5-31. http://dx.doi.org/10.1007/s10584-011-0148-z
- Jacobson, M.Z., 2008: On the causal link between carbon dioxide and air pollution mortality. *Geophysical Research Letters*, 35, L03809. http://dx.doi.org/10.1029/2007GL031101
- Jhun, I., N. Fann, A. Zanobetti, and B. Hubbell, 2014: Effect modification of ozone-related mortality risks by temperature in 97 US cities. *Environment International*, 73, 128-134. http://dx.doi.org/10.1016/j.envint.2014.07.009
- Ren, C., G.M. Williams, K. Mengersen, L. Morawska, and S. Tong, 2008: Does temperature modify short-term effects of ozone on total mortality in 60 large eastern US communities? An assessment using the NMMAPS data. *Environment International*, 34, 451-458. http://dx.doi.org/10.1016/j. envint.2007.10.001
- Ren, C., G.M. Williams, L. Morawska, K. Mengersen, and S. Tong, 2008: Ozone modifies associations between temperature and cardiovascular mortality: Analysis of the NMMAPS data. *Occupational and Environmental Medicine*, 65, 255-260. http://dx.doi.org/10.1136/oem.2007.033878
- Ren, C., G.M. Williams, K. Mengersen, L. Morawska, and S. Tong, 2009: Temperature enhanced effects of ozone on cardiovascular mortality in 95 large US communities, 1987-2000: Assessment using the NMMAPS data. Archives of Environmental & Occupational Health, 64, 177-184. http:// dx.doi.org/10.1080/19338240903240749
- Ren, C., S. Melly, and J. Schwartz, 2010: Modifiers of short-term effects of ozone on mortality in eastern Massachusetts-A casecrossover analysis at individual level. *Environmental Health*, 9, Article 3. http://dx.doi.org/10.1186/1476-069X-9-3
- 56. EPA, 2015: Technical Support Document (TSD): Preparation of Emissions Inventories for the Version 6.2, 2011 Emissions Modeling Platform. U.S. Environmental Protection Agency, Office of Air and Radiation. http://www3.epa.gov/ttn/chief/emch/2011v6/2011v6_2_2017_2025_Emis-Mod_TSD_aug2015.pdf
- Otte, T.L., C.G. Nolte, M.J. Otte, and J.H. Bowden, 2012: Does nudging squelch the extremes in regional climate modeling? *Journal of Climate*, 25, 7046-7066. http://dx.doi. org/10.1175/JCLI-D-12-00048.1

- Gao, Y., J.S. Fu, J.B. Drake, J.-F. Lamarque, and Y. Liu, 2013: The impact of emission and climate change on ozone in the United States under representative concentration pathways (RCPs). *Atmospheric Chemistry and Physics*, 13, 9607-9621. http://dx.doi.org/10.5194/acp-13-9607-2013
- 59. Bell, M.L. and F. Dominici, 2008: Effect modification by community characteristics on the short-term effects of ozone exposure and mortality in 98 US communities. *American Journal of Epidemiology*, **167**, 986-97. http://dx.doi.org/10.1093/aje/kwm396
- Sacks, J.D., L.W. Stanek, T.J. Luben, D.O. Johns, B.J. Buckley, J.S. Brown, and M. Ross, 2011: Particulate matter-induced health effects: Who is susceptible? *Environmental Health Perspectives*, 119, 446-454. http://dx.doi. org/10.1289/ehp.1002255
- EPA, 1999: Regulatory Impact Analysis Control of Air Pollution from New Motor Vehicles: Tier 2 Motor Vehicle Emissions Standards and Gasoline Sulfur Control Requirements. EPA420-R-99-023, 522 pp. U.S. Environmental Protection Agency, Office of Transportation and Air Quality, Washington, D.C. http://www.epa.gov/tier2/documents/ r99023.pdf
- 62. EPA, 2015: Cross-State Air Pollution Rule (CSAPR). U.S. Environmental Protection Agency, Washington, D.C. http:// www.epa.gov/airtransport/CSAPR/
- 63. EPA, 2015: Tier 2 Vehicle and Gasoline Sulfur Program. U.S. Environmental Protection Agency, Office of Transportation and Air Quality, Washington, D.C. http://www.epa. gov/tier2/
- Dawson, J.P., P.N. Racherla, B.H. Lynn, P.J. Adams, and S.N. Pandis, 2009: Impacts of climate change on regional and urban air quality in the eastern United States: Role of meteorology. *Journal of Geophysical Research: Atmospheres*, 114. http://dx.doi.org/10.1029/2008JD009849
- 65. Trail, M., A.P. Tsimpidi, P. Liu, K. Tsigaridis, J. Rudokas, P. Miller, A. Nenes, Y. Hu, and A.G. Russell, 2014: Sensitivity of air quality to potential future climate change and emissions in the United States and major cities. *Atmospheric Environment*, **94**, 552-563. http://dx.doi.org/10.1016/j. atmosenv.2014.05.079
- 66. Delfino, R.J., S. Brummel, J. Wu, H. Stern, B. Ostro, M. Lipsett, A. Winer, D.H. Street, L. Zhang, T. Tjoa, and D.L. Gillen, 2009: The relationship of respiratory and cardiovas-cular hospital admissions to the southern California wildfires of 2003. *Occupational and Environmental Medicine*, 66, 189-197. http://dx.doi.org/10.1136/oem.2008.041376

- Künzli, N., E. Avol, J. Wu, W.J. Gauderman, E. Rappaport, J. Millstein, J. Bennion, R. McConnell, F.D. Gilliland, K. Berhane, F. Lurmann, A. Winer, and J.M. Peters, 2006: Health effects of the 2003 southern California wildfires on children. *American Journal of Respiratory and Critical Care Medicine*, **174**, 1221-1228. http://dx.doi.org/10.1164/ rccm.200604-519OC
- Park, R.J., 2003: Sources of carbonaceous aerosols over the United States and implications for natural visibility. *Journal of Geophysical Research*, **108**, D12, 4355. http://dx.doi. org/10.1029/2002JD003190
- Sapkota, A., J.M. Symons, J. Kleissl, L. Wang, M.B. Parlange, J. Ondov, P.N. Breysse, G.B. Diette, P.A. Eggleston, and T.J. Buckley, 2005: Impact of the 2002 Canadian forest fires on particulate matter air quality in Baltimore City. *Environmental Science & Technology*, **39**, 24-32. http://dx.doi.org/10.1021/es035311z
- Henderson, S.B., M. Brauer, Y.C. Macnab, and S.M. Kennedy, 2011: Three measures of forest fire smoke exposure and their associations with respiratory and cardiovascular health outcomes in a population-based cohort. *Environmental Health Perspectives*, **119**, 1266-1271. http://dx.doi.org/10.1289/ehp.1002288
- Rappold, A.G., W.E. Cascio, V.J. Kilaru, S.L. Stone, L.M. Neas, R.B. Devlin, and D. Diaz-Sanchez, 2012: Cardio-respiratory outcomes associated with exposure to wildfire smoke are modified by measures of community health. *Environmental Health*, **11**, Article 71. http://dx.doi. org/10.1186/1476-069X-11-71
- 72. Liu, J.C., G. Pereira, S.A. Uhl, M.A. Bravo, and M.L. Bell, 2015: A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environmental Research*, **136**, 120-132. http://dx.doi.org/10.1016/j. envres.2014.10.015
- Westerling, A.L., H.G. Hidalgo, D.R. Cayan, and T.W. Swetnam, 2006: Warming and earlier spring increase western U.S. forest wildfire activity. *Science*, 313, 940-943. http:// dx.doi.org/10.1126/science.1128834
- 74. Walsh, J., D. Wuebbles, K. Hayhoe, J. Kossin, K. Kunkel, G. Stephens, P. Thorne, R. Vose, M. Wehner, J. Willis, D. Anderson, S. Doney, R. Feely, P. Hennon, V. Kharin, T. Knutson, F. Landerer, T. Lenton, J. Kennedy, and R. Somerville, 2014: Ch. 2: Our changing climate. *Climate Change Impacts in the United States: The Third National Climate Assessment*. Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds. U.S. Global Change Research Program, Washington, D.C., 19-67. http://dx.doi.org/10.7930/J0KW5CXT

- 75. Garfin, G., G. Franco, H. Blanco, A. Comrie, P. Gonzalez, T. Piechota, R. Smyth, and R. Waskom, 2014: Ch. 20: Southwest. *Climate Change Impacts in the United States: The Third National Climate Assessment*. Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds. U.S. Global Change Research Program, Washington, D.C., 462-486. http://dx.doi.org/10.7930/J08G8HMN
- 76. Luber, G., K. Knowlton, J. Balbus, H. Frumkin, M. Hayden, J. Hess, M. McGeehin, N. Sheats, L. Backer, C.B. Beard, K.L. Ebi, E. Maibach, R.S. Ostfeld, C. Wiedinmyer, E. Zielinski-Gutiérrez, and L. Ziska, 2014: Ch. 9: Human health. *Climate Change Impacts in the United States: The Third National Climate Assessment*. Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds. U.S. Global Change Research Program, Washington, D.C., 220-256. http://dx.doi.org/10.7930/J0PN93H5
- Westerling, A.L., M.G. Turner, E.A.H. Smithwick, W.H. Romme, and M.G. Ryan, 2011: Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. *Proceedings of the National Academy of Sciences*, 108, 13165-13170. http://dx.doi.org/10.1073/pnas.1110199108
- 78. Keywood, M., M. Kanakidou, A. Stohl, F. Dentener, G. Grassi, C.P. Meyer, K. Torseth, D. Edwards, A.M. Thompson, U. Lohmann, and J. Burrows, 2013: Fire in the air: Biomass burning impacts in a changing climate. *Critical Reviews in Environmental Science and Technology*, 43, 40-83. http://dx.doi.org/10.1080/10643389.2011.604248
- Spracklen, D.V., L.J. Mickley, J.A. Logan, R.C. Hudman, R. Yevich, M.D. Flannigan, and A.L. Westerling, 2009: Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous aerosol concentrations in the western United States. *Journal of Geophysical Research: Atmospheres*, **114**, D20301. http://dx.doi.org/10.1029/2008JD010966
- Seager, R., M. Ting, I. Held, Y. Kushnir, J. Lu, G. Vecchi, H.-P. Huang, N. Harnik, A. Leetmaa, N.-C. Lau, C. Li, J. Velez, and N. Naik, 2007: Model projections of an imminent transition to a more arid climate in southwestern North America. *Science*, **316**, 1181-1184. http://dx.doi.org/10.1126/science.1139601
- Reid, C.E. and J.L. Gamble, 2009: Aeroallergens, allergic disease, and climate change: Impacts and adaptation. *Eco-Health*, 6, 458-470. http://dx.doi.org/10.1007/s10393-009-0261-x

- Selgrade, M.K., R.F. Lemanske, Jr., M.I. Gilmour, L.M. Neas, M.D.W. Ward, P.K. Henneberger, D.N. Weissman, J.A. Hoppin, R.R. Dietert, P.D. Sly, A.M. Geller, P.L. Enright, G.S. Backus, P.A. Bromberg, D.R. Germolec, and K.B. Yeatts, 2006: Induction of asthma and the environment: What we know and need to know. *Environmental Health Perspectives*, **114**, 615-619. http://dx.doi.org/10.1289/ ehp.8376
- Kinney, P.L., 2008: Climate change, air quality, and human health. *American Journal of Preventive Medicine*, **35**, 459-467. http://dx.doi.org/10.1016/j.amepre.2008.08.025
- Bielory, L., K. Lyons, and R. Goldberg, 2012: Climate change and allergic disease. *Current Allergy and Asthma Reports*, 12, 485-494. http://dx.doi.org/10.1007/s11882-012-0314-z
- 85. AAAAI, Undated: Allergy Statistics. American Academy of Allergy, Asthma and Immunology. http://www.aaaai.org/ about-the-aaaai/newsroom/allergy-statistics.aspx
- Breton, M.-C., M. Garneau, I. Fortier, F. Guay, and J. Louis, 2006: Relationship between climate, pollen concentrations of *Ambrosia* and medical consultations for allergic rhinitis in Montreal, 1994–2002. *Science of The Total Environment*, **370**, 39-50. http://dx.doi.org/10.1016/j.scitotenv.2006.05.022
- Bartra, J., J. Mullol, A. del Cuvillo, I. Davila, M. Ferrer, I. Jauregui, J. Montoro, J. Sastre, and A. Valero, 2007: Air pollution and allergens. *Journal of Investigational Allergology and Clinical Immunology*, **17 Suppl 2**, 3-8. http://www.jiaci.org/ issues/vol17s2/vol17s2-2.htm
- Blando, J., L. Bielory, V. Nguyen, R. Diaz, and H.A. Jeng, 2012: Anthropogenic climate change and allergic diseases. *Atmosphere*, **3**, 200-212. http://dx.doi.org/10.3390/ atmos3010200
- EPA, 2008: Review of the Impacts of Climate Variability and Change on Aeroallergens and Their Associated Effects. EPA/600/R-06/164F, 125 pp. U.S. Environmental Protection Agency, Washington, D.C. http://ofmpub.epa.gov/ eims/eimscomm.getfile?p_download_id=490474
- Lawson, J.A. and A. Senthilselvan, 2005: Asthma epidemiology: Has the crisis passed? *Current Opinion in Pulmonary Medicine*, 11, 79-84.
- 91. Beggs, P.J., 2004: Impacts of climate change on aeroallergens: Past and future. *Clinical & Experimental Aller*gy, 34, 1507-1513. http://dx.doi.org/10.1111/j.1365-2222.2004.02061.x

- 92. Beggs, P.J. and H.J. Bambrick, 2005: Is the global rise of asthma an early impact of anthropogenic climate change? *Environmental Health Perspectives*, **113**, 915-919. http://dx. doi.org/10.1289/ehp.7724
- 93. D'Amato, G., C.E. Baena-Cagnani, L. Cecchi, I. Annesi-Maesano, C. Nunes, I. Ansotegui, M. D'Amato, G. Liccardi, M. Sofia, and W.G. Canonica, 2013: Climate change, air pollution and extreme events leading to increasing prevalence of allergic respiratory diseases. *Multidisciplinary Respiratory Medicine*, **8**, 1-9. http://dx.doi.org/10.1186/2049-6958-8-12
- Albertine, J.M., W.J. Manning, M. DaCosta, K.A. Stinson, M.L. Muilenberg, and C.A. Rogers, 2014: Projected carbon dioxide to increase grass pollen and allergen exposure despite higher ozone levels. *PLoS ONE*, 9, e111712. http://dx.doi. org/10.1371/journal.pone.0111712
- 95. Ziska, L., K. Knowlton, C. Rogers, D. Dalan, N. Tierney, M.A. Elder, W. Filley, J. Shropshire, L.B. Ford, C. Hedberg, P. Fleetwood, K.T. Hovanky, T. Kavanaugh, G. Fulford, R.F. Vrtis, J.A. Patz, J. Portnoy, F. Coates, L. Bielory, and D. Frenz, 2011: Recent warming by latitude associated with increased length of ragweed pollen season in central North America. *Proceedings of the National Academy of Sciences*, **108**, 4248-4251. http://dx.doi.org/10.1073/pnas.1014107108
- 96. Rogers, C.A., P.M. Wayne, E.A. Macklin, M.L. Muilenberg, C.J. Wagner, P.R. Epstein, and F.A. Bazzaz, 2006: Interaction of the onset of spring and elevated atmospheric CO₂ on ragweed (*Ambrosia artemisiifolia* L.) pollen production. *Environmental Health Perspectives*, **114**, 865-869. http://dx. doi.org/10.1289/ehp.8549
- 97. Singer, B.D., L.H. Ziska, D.A. Frenz, D.E. Gebhard, and J.G. Straka, 2005: Increasing Amb a 1 content in common ragweed (*Ambrosia artemisiifolia*) pollen as a function of rising atmospheric CO₂ concentration. *Functional Plant Biology*, **32**, 667-670. http://dx.doi.org/10.1071/fp05039
- Neil, K. and J. Wu, 2006: Effects of urbanization on plant flowering phenology: A review. Urban Ecosystems, 9, 243-257. http://dx.doi.org/10.1007/s11252-006-9354-2
- George, K., L.H. Ziska, J.A. Bunce, and B. Quebedeaux, 2007: Elevated atmospheric CO₂ concentration and temperature across an urban–rural transect. *Atmospheric Environment*, **41**, 7654-7665. http://dx.doi.org/10.1016/j. atmosenv.2007.08.018

- 100. Roetzer, T., M. Wittenzeller, H. Haeckel, and J. Nekovar, 2000: Phenology in central Europe: Differences and trends of spring phenophases in urban and rural areas. *International Journal of Biometeorology*, **44**, 60-66. http://dx.doi. org/10.1007/s004840000062
- 101. Garcia-Mozo, H., C. Galán, V. Jato, J. Belmonte, C.D. de la Guardia, D. Fernández, M. Gutiérrez, M.J. Aira, J.M. Roure, L. Ruiz, M.M. Trigo, and E. Domínguez-Vilches, 2006: *Quercus* pollen season dynamics in the Iberian peninsula: Response to meteorological parameters and possible consequences of climate change. *Annals of Agricultural and Environmental Medicine*, **13**, 209-224. http://www.uco. es/aerobiologia/publicaciones/modelling/climate_change/ Quercus_AAEM_def.pdf
- LaDeau, S.L. and J.S. Clark, 2006: Elevated CO₂ and tree fecundity: The role of tree size, interannual variability, and population heterogeneity. *Global Change Biology*, **12**, 822-833. http://dx.doi.org/10.1111/j.1365-2486.2006.01137.x
- 103. Zhang, R., T. Duhl, M.T. Salam, J.M. House, R.C. Flagan, E.L. Avol, F.D. Gilliland, A. Guenther, S.H. Chung, B.K. Lamb, and T.M. VanReken, 2013: Development of a regional-scale pollen emission and transport modeling framework for investigating the impact of climate change on allergic airway disease. *Biogeosciences*, **10**, 3977-4023. http://dx.doi. org/10.5194/bgd-10-3977-2013
- 104. D'Amato, G. and L. Cecchi, 2008: Effects of climate change on environmental factors in respiratory allergic diseases. *Clinical & Experimental Allergy*, **38**, 1264-1274. http://dx. doi.org/10.1111/j.1365-2222.2008.03033.x
- 105. D'Amato, G., M. Rottem, R. Dahl, M.S. Blaiss, E. Ridolo, L. Cecchi, N. Rosario, C. Motala, I. Ansotegui, and I. Annesi-Maesano, 2011: Climate change, migration, and allergic respiratory diseases: An update for the allergist. *World Allergy Organization Journal*, 4, 121-125. http://dx. doi.org/10.1097/WOX.0b013e3182260a57
- 106. Shea, K.M., R.T. Truckner, R.W. Weber, and D.B. Peden, 2008: Climate change and allergic disease. *Journal of Aller*gy and Clinical Immunology, **122**, 443-453. http://dx.doi. org/10.1016/j.jaci.2008.06.032
- 107. D'Amato, G., L. Cecchi, M. D'Amato, and G. Liccardi, 2010: Urban air pollution and climate change as environmental risk factors of respiratory allergy: An update. *Journal* of *Investigational Allergology and Clinical Immunology*, 20, 95-102. http://www.jiaci.org/issues/vol20issue2/1.pdf

- Cecchi, L., G. D'Amato, J.G. Ayres, C. Galan, F. Forastiere, B. Forsberg, J. Gerritsen, C. Nunes, H. Behrendt, C. Akdis, R. Dahl, and I. Annesi-Maesano, 2010: Projections of the effects of climate change on allergic asthma: The contribution of aerobiology. *Allergy*, 65, 1073-1081. http://dx.doi. org/10.1111/j.1398-9995.2010.02423.x
- 109. D'Amato, G., G. Liccardi, and G. Frenguelli, 2007: Thunderstorm-asthma and pollen allergy. *Allergy*, **62**, 11-16. http://dx.doi.org/10.1111/j.1398-9995.2006.01271.x
- 110. D'Amato, G., G. Liccardi, M. D'Amato, and M. Cazzola, 2001: The role of outdoor air pollution and climatic changes on the rising trends in respiratory allergy. *Respiratory Medicine*, **95**, 606-611. http://dx.doi.org/10.1053/ rmed.2001.1112
- 111. D'Amato, G., 2002: Environmental urban factors (air pollution and allergens) and the rising trends in allergic respiratory diseases. *Allergy*, **57**, 30-33. http://dx.doi. org/10.1034/j.1398-9995.57.s72.5.x
- 112. D'Amato, G., G. Liccardi, M. D'Amato, and S. Holgate, 2005: Environmental risk factors and allergic bronchial asthma. *Clinical & Experimental Allergy*, **35**, 1113-1124. http:// dx.doi.org/10.1111/j.1365-2222.2005.02328.x
- 113. Atkinson, R.W. and D.P. Strachan, 2004: Role of outdoor aeroallergens in asthma exacerbations: Epidemiological evidence. *Thorax*, **59**, 277-278. http://dx.doi.org/10.1136/ thx.2003.019133
- 114. Zhong, W., L. Levin, T. Reponen, G.K. Hershey, A. Adhikari, R. Shukla, and G. LeMasters, 2006: Analysis of short-term influences of ambient aeroallergens on pediatric asthma hospital visits. *Science of The Total Environment*, **370**, 330-336. http://dx.doi.org/10.1016/j.scitotenv.2006.06.019
- 115. Cakmak, S., R.E. Dales, and F. Coates, 2012: Does air pollution increase the effect of aeroallergens on hospitalization for asthma? *Journal of Allergy and Clinical Immunology*, **129**, 228-231. http://dx.doi.org/10.1016/j.jaci.2011.09.025
- 116. IOM, 2011: Climate Change, the Indoor Environment, and Health. Institute of Medicine. The National Academies Press, Washington, D.C., 286 pp. http://dx.doi. org/10.17226/13115
- 117. IOM, 1993: Indoor Allergens: Assessing and Controlling Adverse Health Effects. Institute of Medicine. The National Academies Press, Washington, D.C., 350 pp. http://dx.doi. org/10.17226/2056

- 118. IOM, 2000: Clearing the Air: Asthma and Indoor Air Exposures. Institute of Medicine. The National Academies Press, Washington, D.C., 456 pp. http://www.nap.edu/catalog/9610/clearing-the-air-asthma-and-indoor-air-exposures
- IOM, 2004: Damp Indoor Spaces and Health. Institute of Medicine. The National Academies Press, Washington, D.C., 370 pp. http://dx.doi.org/10.17226/11011
- 120. WHO, 2009: WHO Handbook on Indoor Radon: A Public Health Perspective. Zeeb, H. and F. Shannoun, Eds. World Health Organization, Geneva, Switzerland, 108 pp. http:// whqlibdoc.who.int/publications/2009/9789241547673_ eng.pdf
- 121. WHO, 2010: WHO Guidelines for Indoor Air Quality: Selected Pollutants. 484 pp. World Health Organization, Geneva. http://www.euro.who.int/__data/assets/pdf_ file/0009/128169/e94535.pdf
- 122. Ott, W.R., 1989: Human activity patterns: A review of the literature for estimating time spent indoors, outdoors, and in transit. *Proceedings of the Research Planning Conference on Human Activity Patterns*. EPA National Exposure Research Laboratory, Las Vegas, NV, 3-1 to 3-38.
- 123. Klepeis, N.E., W.C. Nelson, W.R. Ott, J.P. Robinson, A.M. Tsang, P. Switzer, J.V. Behar, S.C. Hern, and W.H. Engelmann, 2001: The National Human Activity Pattern Survey (NHAPS): A resource for assessing exposure to environmental pollutants. *Journal of Exposure Analysis and Environmental Epidemiology*, **11**, 231-252. http://dx.doi.org/10.1038/ sj.jea.7500165
- 124. Riley, W.J., T.E. McKone, A.C. Lai, and W.W. Nazaroff, 2002: Indoor particulate matter of outdoor origin: Importance of size-dependent removal mechanisms. *Environmental Science & Technology*, **36**, 200-207. http://dx.doi. org/10.1021/es010723y
- 125. Bennett, D.H. and P. Koutrakis, 2006: Determining the infiltration of outdoor particles in the indoor environment using a dynamic model. *Journal of Aerosol Science*, **37**, 766-785. http://dx.doi.org/10.1016/j.jaerosci.2005.05.020
- 126. Wallace, L., 1996: Indoor particles: A review. Journal of the Air & Waste Management Association, 46, 98-126. http:// dx.doi.org/10.1080/10473289.1996.10467451
- 127. Brennan, T., J.B. Cummings, and J. Lstiburek, 2002: Unplanned airflows & moisture problems. *ASHRAE journal*, 44, 44-49. https://64.94.228.53/File Library/docLib/Public/2002102583440_266.pdf

- 128. Stephens, B. and J.A. Siegel, 2012: Penetration of ambient submicron particles into single-family residences and associations with building characteristics. *Indoor Air*, **22**, 501-513. http://dx.doi.org/10.1111/j.1600-0668.2012.00779.x
- Persily, A., A. Musser, and S.J. Emmerich, 2010: Modeled infiltration rate distributions for U.S. housing. *Indoor Air*, 20, 473-485. http://dx.doi.org/10.1111/j.1600-0668.2010.00669.x
- 130. Fisk, W.J., 2015: Review of some effects of climate change on indoor environmental quality and health and associated no-regrets mitigation measures. *Building and Environment*, 86, 70-80. http://dx.doi.org/10.1016/j.buildenv.2014.12.024
- 131. Ilacqua, V., J. Dawson, M. Breen, S. Singer, and A. Berg, 2015: Effects of climate change on residential infiltration and air pollution exposure. *Journal of Exposure Science and Environmental Epidemiology*, Published online 27 May 2015. http://dx.doi.org/10.1038/jes.2015.38
- Weschler, C.J., 2006: Ozone's impact on public health: Contributions from indoor exposures to ozone and products of ozone-initiated chemistry. *Environmental Health Perspectives*, 114, 1489-1496. PMC1626413
- 133. Bell, M.L., A. Zanobetti, and F. Dominici, 2014: Who is more affected by ozone pollution? A systematic review and meta-analysis. *American Journal of Epidemiology*, 180, 15-28. http://dx.doi.org/10.1093/aje/kwu115
- 134. Lin, S., X. Liu, L.H. Le, and S.-A. Hwang, 2008: Chronic exposure to ambient ozone and asthma hospital admissions among children. *Environmental Health Perspectives*, **116**, 1725-1730. http://dx.doi.org/10.1289/ehp.11184
- 135. Nazaroff, W.W., 2013: Exploring the consequences of climate change for indoor air quality. *Environmental Research Letters*,
 8. http://dx.doi.org/10.1088/1748-9326/8/1/015022
- 136. Bowers, R.M., N. Clements, J.B. Emerson, C. Wiedinmyer, M.P. Hannigan, and N. Fierer, 2013: Seasonal variability in bacterial and fungal diversity of the near-surface atmospher. *Environmental Science & Technology*, 47, 12097-12106. http://dx.doi.org/10.1021/es402970s
- 137. Hardin, B.D., B.J. Kelman, and A. Saxon, 2003: Adverse human health effects associated with molds in the indoor environment. *Journal of Occupational and Environmental Medicine*, **45**, 470-478. http://dx.doi.org/10.1097/00043764-200305000-00006

- 138. Walzer, P.D., 2013: The ecology of pneumocystis: Perspectives, personal recollections, and future research opportunities. *The Journal of Eukaryotic Microbiology*, **60**, 634-645. http://dx.doi.org/10.1111/jeu.12072
- 139. Parr, A., E.A. Whitney, and R.L. Berkelman, 2014: Legionellosis on the rise: A review of guidelines for prevention in the United States. *Journal of Public Health Management and Practice*, **21**, E17-E26. http://dx.doi.org/10.1097/ phh.00000000000123
- Decker, B.K. and T.N. Palmore, 2014: Hospital water and opportunities for infection prevention. *Curr Infect Dis Rep*, 16, 432. http://dx.doi.org/10.1007/s11908-014-0432-y
- 141. Cunha, B.A., A. Burillo, and E. Bouza, 2015: Legionnaires' disease. *Lancet*, Published online 28 July 2015. http://dx. doi.org/10.1016/S0140-6736(15)60078-2
- 142. Hines, S.A., D.J. Chappie, R.A. Lordo, B.D. Miller, R.J. Janke, H.A. Lindquist, K.R. Fox, H.S. Ernst, and S.C. Taft, 2014: Assessment of relative potential for *Legionella* species or surrogates inhalation exposure from common water uses. *Water Research*, **56**, 203-213. http://dx.doi.org/10.1016/j. watres.2014.02.013
- 143. Whiley, H., S. Giglio, and R. Bentham, 2015: Opportunistic pathogens Mycobacterium Avium Complex (MAC) and *Legionella* spp. colonise model shower *Pathogens*, 4, 590-598. http://dx.doi.org/10.3390/pathogens4030590
- 144. Phin, N., F. Parry-Ford, T. Harrison, H.R. Stagg, N. Zhang, K. Kumar, O. Lortholary, A. Zumla, and I. Abubakar, 2014: Epidemiology and clinical management of Legionnaires' disease. *The Lancet Infectious Diseases*, 14, 1011-1021. http:// dx.doi.org/10.1016/s1473-3099(14)70713-3
- 145. Falkinham, J.O., III,, E.D. Hilborn, M.J. Arduino, A. Pruden, and M.A. Edwards, 2015: Epidemiology and ecology of opportunistic premise plumbing pathogens: *Legionella pneumophila*, *Mycobacterium avium*, and *Pseudomonas aeruginosa*. *Environmental Health Perspectives*, **123**, 749-758. http://dx. doi.org/10.1289/ehp.1408692
- 146. Halsby, K.D., C.A. Joseph, J.V. Lee, and P. Wilkinson, 2014: The relationship between meteorological variables and sporadic cases of Legionnaires' disease in residents of England and Wales. *Epidemiology and Infection*, **142**, 2352-2359. http://dx.doi.org/10.1017/S0950268813003294
- 147. Cunha, B.A., J. Connolly, and E. Abruzzo, 2015: Increase in pre-seasonal community-acquired Legionnaire's disease due to increased precipitation. *Clinical Microbiology* and Infection, 21, e45-e46. http://dx.doi.org/10.1016/j. cmi.2015.02.015

- Hicks, L.A., L.E. Garrison, G.E. Nelson, and L.M. Hampton, 2011: Legionellosis---United States, 2000-2009. MMWR. Morbidity and Mortality Weekly Report, 60, 1083-1086. http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6032a3.htm
- 149. Farnham, A., L. Alleyne, D. Cimini, and S. Balter, 2014: Legionnaires' disease incidence and risk factors, New York, New York, USA, 2001-2011. *Emerging Infectious Diseases*, 20, 1795-1802. http://dx.doi.org/10.3201/eid2011.131872
- 150. Johanning, E., P. Auger, P.R. Morey, C.S. Yang, and E. Olmsted, 2014: Review of health hazards and prevention measures for response and recovery workers and volunteers after natural disasters, flooding, and water damage: Mold and dampness. *Environmental Health and Preventive Medicine*, 19, 93-99. http://dx.doi.org/10.1007/s12199-013-0368-0
- 151. Mudarri, D. and W.J. Fisk, 2007: Public health and economic impact of dampness and mold. *Indoor Air*, **17**, 226-235. http://dx.doi.org/10.1111/j.1600-0668.2007.00474.x
- 152. Fisk, W.J., E.A. Eliseeva, and M.J. Mendell, 2010: Association of residential dampness and mold with respiratory tract infections and bronchitis: A meta-analysis. *Environmental Health*, **9**, Article 72. http://dx.doi.org/10.1186/1476-069x-9-72
- Seltenrich, N., 2012: Healthier tribal housing: Combining the best of old and new. *Environmental Health Perspectives*, 120, A460-A469. http://dx.doi.org/10.1289/ehp.120-a460
- 154. Parthasarathy, S., R.L. Maddalena, M.L. Russell, and M.G. Apte, 2011: Effect of temperature and humidity on formaldehyde emissions in temporary housing units. *Journal of the Air & Waste Management Association*, **61**, 689-695. http:// dx.doi.org/10.3155/1047-3289.61.6.689
- 155. Norbäck, D., G. Wieslander, K. Nordström, and R. Wålinder, 2000: Asthma symptoms in relation to measured building dampness in upper concrete floor construction, and 2-ethyl-1-hexanol in indoor air. *International Journal of Tuberculosis and Lung Disease*, 4, 1016-1025. http://www.nchh.org/portals/0/contents/article0877.pdf
- 156. Markowicz, P. and L. Larsson, 2015: Influence of relative humidity on VOC concentrations in indoor air. *Environmental Science and Pollution Research*, **22**, 5772-5779. http://dx.doi.org/10.1007/s11356-014-3678-x

- 157. Waite, T., V. Murray, and D. Baker, 2014: Carbon monoxide poisoning and flooding: Changes in risk before, during and after flooding require appropriate public health interventions. *Plos Currents Disasters*, July 3, Edition 1. http://dx.doi.org/10.1371/currents.dis.2b2eb9e-15f9b982784938803584487f1
- 158. Bronstein, A., J.H. Clower, S. Iqbal, F.Y. Yip, C.A. Martin, A. Chang, A.F. Wolkin, and J. Bell, 2011: Carbon monoxide exposures---United States, 2000--2009. *MMWR. Morbidity* and Mortality Weekly Report, **60**, 1014-1017. http://www. cdc.gov/mmwr/preview/mmwrhtml/mm6030a2.htm
- 159. Watson, D.C., M. Sargianou, A. Papa, P. Chra, I. Starakis, and G. Panos, 2014: Epidemiology of Hantavirus infections in humans: A comprehensive, global overview. *Critical Reviews in Microbiology*, 40, 261-272. http://dx.doi.org/10. 3109/1040841x.2013.783555
- 160. Jonsson, C.B., L.T.M. Figueiredo, and O. Vapalahti, 2010: A global perspective on hantavirus ecology, epidemiology, and disease. *Clinical Microbiology Reviews*, 23, 412-441. http://dx.doi.org/10.1128/cmr.00062-09
- 161. Dearing, M.D. and L. Dizney, 2010: Ecology of hantavirus in a changing world. *Annals of the New York Academy of Sciences*, **1195**, 99-112. http://dx.doi.org/10.1111/j.1749-6632.2010.05452.x
- 162. Bezirtzoglou, C., K. Dekas, and E. Charvalos, 2011: Climate changes, environment and infection: Facts, scenarios and growing awareness from the public health community within Europe. *Anaerobe*, **17**, 337-340. http://dx.doi. org/10.1016/j.anaerobe.2011.05.016
- 163. Klein, S.L. and C.H. Calisher, 2007: Emergence and persistence of hantaviruses. Wildlife and Emerging Zoonotic Diseases: The Biology, Circumstances and Consequences of Cross-Species Transmission. Childs, J.E., J.S. Mackenzie, and J.A. Richt, Eds. Springer-Verlag, Berlin, 217-252. http:// dx.doi.org/10.1007/978-3-540-70962-6_10
- Reusken, C. and P. Heyman, 2013: Factors driving hantavirus emergence in Europe. *Current Opinion in Virology*, 3, 92-99. http://dx.doi.org/10.1016/j.coviro.2013.01.002
- 165. Medina-Ramón, M. and J. Schwartz, 2007: Temperature, temperature extremes, and mortality: A study of acclimatisation and effect modification in 50 US cities. *Occupational* and Environmental Medicine, 64, 827-833. http://dx.doi. org/10.1136/oem.2007.033175

- 166. Akinbami, L.J., J.E. Moorman, C. Bailey, H.S. Zahran, M. King, C.A. Johnson, and X. Liu, 2012: Trends in Asthma Prevalence, Health Care Use, and Mortality in the United States, 2001–2010. NCHS Data Brief No. 94, May 2012, 8 pp. National Center for Health Statistics, Hyattsville, MD. http://www.cdc.gov/nchs/data/databriefs/db94.pdf
- 167. Adamkiewicz, G., A.R. Zota, P. Fabian, T. Chahine, R. Julien, J.D. Spengler, and J.I. Levy, 2011: Moving environmental justice indoors: Understanding structural influences on residential exposure patterns in low-income communities. *American Journal of Public Health*, **101**, S238-S245. http://dx.doi.org/10.2105/AJPH.2011.300119
- 168. Kitch, B.T., G. Chew, H.A. Burge, M.L. Muilenberg, S.T. Weiss, T.A. Platts-Mills, G. O'Connor, and D.R. Gold, 2000: Socioeconomic predictors of high allergen levels in homes in the greater Boston area. *Environmental Health Perspectives*, **108**, 301-307. PMC1638021
- 169. Bloom, B., L.I. Jones, and G. Freeman, 2013: Summary Health Statistics for U.S. Children: National Health Interview Survey, 2012. Vital and Health Statistics 10(258), 73 pp. National Center for Health Statistics, Hyattsville, MD. http://www.cdc.gov/nchs/data/series/sr_10/sr10_258.pdf
- 170. Gelfand, E.W., 2009: Pediatric asthma: A different disease. *Proceedings of the American Thoracic Society*, **6**, 278-282. http://dx.doi.org/10.1513/pats.200808-090RM
- 171. DellaValle, C.T., E.W. Triche, B.P. Leaderer, and M.L. Bell, 2012: Effects of ambient pollen concentrations on frequency and severity of asthma symptoms among asthmatic children. *Epidemiology*, 23, 55-63. http://dx.doi.org/10.1097/ EDE.0b013e31823b66b8
- 172. Ebi, K.L. and J.A. Paulson, 2007: Climate change and children. *Pediatric Clinics of North America*, **54**, 213-226. http://dx.doi.org/10.1016/j.pcl.2007.01.004
- 173. Schmier, J.K. and K.L. Ebi, 2009: The impact of climate change and aeroallergens on children's health. *Allergy and Asthma Proceedings*, **30**, 229-237. http://dx.doi. org/10.2500/aap.2009.30.3229
- 174. Moorman, J.E., L.J. Akinbami, C.M. Bailey, H.S. Zahran, M.E. King, C.A. Johnson, and X. Liu, 2012: National Surveillance of Asthma: United States, 2001-2010. Vital and Health Statistics 3(35). National Center for Health Statistics. http://www.cdc.gov/nchs/data/series/sr_03/sr03_035. pdf

- 175. Meng, Y.Y., S.H. Babey, T.A. Hastert, and E.R. Brown, 2007: California's racial and ethnic minorities more adversely affected by asthma. *Policy Brief UCLA Center for Health Policy Research* 1-7. https://escholarship.org/uc/item/4k45v3xt
- 176. Brim, S.N., R.A. Rudd, R.H. Funk, and D.B. Callahan, 2008: Asthma prevalence among US children in underrepresented minority populations: American Indian/Alaska Native, Chinese, Filipino, and Asian Indian. *Pediatrics*, **122**, e217-e222. http://dx.doi.org/10.1542/peds.2007-3825
- 177. Crocker, D., C. Brown, R. Moolenaar, J. Moorman, C. Bailey, D. Mannino, and F. Holguin, 2009: Racial and ethnic disparities in asthma medication usage and health-care utilization: Data from the National Asthma Survey. *Chest*, **136**, 1063-1071. http://dx.doi.org/10.1378/chest.09-0013
- 178. Miller, J.E., 2000: The effects of race/ethnicity and income on early childhood asthma prevalence and health care use. *American Journal of Public Health*, **90**, 428-430. Pmc1446167
- 179. Peel, J.L., K.B. Metzger, M. Klein, W.D. Flanders, J.A. Mulholland, and P.E. Tolbert, 2007: Ambient air pollution and cardiovascular emergency department visits in potentially sensitive groups. *American Journal of Epidemiology*, 165, 625-633. http://dx.doi.org/10.1093/aje/kwk051

